

CONDITIONAL UPPER BOUND FOR THE k -TH PRIME IDEAL WITH GIVEN ARTIN SYMBOL

LOIĆ GRENIÉ AND GIUSEPPE MOLTENI

ABSTRACT. We prove an explicit upper bound for the k -th prime ideal with fixed Artin symbol, under the assumption of the validity of the Riemann hypothesis for the Dedekind zeta functions.

1. INTRODUCTION

We recall some definitions, just to fix the notations. Let \mathbb{K} be a number field, let $n_{\mathbb{K}}$ denote its dimension, $\Delta_{\mathbb{K}}$ the absolute value of its discriminant, and $r_1(\mathbb{K})$, $r_2(\mathbb{K})$ the number of its real and complex places, respectively. The von Mangoldt function $\Lambda_{\mathbb{K}}$ is defined on the set of ideals of $\mathcal{O}_{\mathbb{K}}$ as $\Lambda_{\mathbb{K}}(\mathfrak{J}) := \log N\mathfrak{p}$ if $\mathfrak{J} = \mathfrak{p}^m$ for some \mathfrak{p} and $m \geq 1$, and is zero otherwise, where \mathfrak{p} denotes any nonzero prime ideal and $N\mathfrak{p}$ its absolute norm.

Moreover, let $\mathbb{K} \subseteq \mathbb{L}$ be a Galois extension of number fields with relative discriminant $\Delta_{\mathbb{L}/\mathbb{K}}$. For \mathfrak{P} a prime ideal of \mathbb{L} above a non-ramified \mathfrak{p} of \mathbb{K} , the Artin symbol $\left[\frac{\mathbb{L}/\mathbb{K}}{\mathfrak{P}}\right]$ denotes the Frobenius automorphism corresponding to $\mathfrak{P}/\mathfrak{p}$, and $\left[\frac{\mathbb{L}/\mathbb{K}}{\mathfrak{p}}\right]$ the conjugacy class of all the $\left[\frac{\mathbb{L}/\mathbb{K}}{\mathfrak{P}}\right]$. The symbol $\left[\frac{\mathbb{L}/\mathbb{K}}{\cdot}\right]$ is then extended multiplicatively to the group of fractional ideals of \mathbb{K} coprime to $\Delta_{\mathbb{L}/\mathbb{K}}$.

Finally, let C be any conjugacy class in $G := \text{Gal}(\mathbb{L}/\mathbb{K})$ and let ε_C be its characteristic function. Then the function π_C and the Chebyshev function ψ_C are defined as

$$\begin{aligned} \pi_C(x) &:= \#\left\{\mathfrak{p} : \mathfrak{p} \text{ non-ramified in } \mathbb{L}/\mathbb{K}, N\mathfrak{p} \leq x, \left[\frac{\mathbb{L}/\mathbb{K}}{\mathfrak{p}}\right] = C\right\} \\ &= \sum_{\substack{\mathfrak{p} \text{ non-ram.} \\ N\mathfrak{p} \leq x}} \varepsilon_C\left(\left[\frac{\mathbb{L}/\mathbb{K}}{\mathfrak{p}}\right]\right), \\ \psi_C(x) &:= \sum_{\substack{\mathfrak{J} \subset \mathcal{O}_{\mathbb{K}} \\ \mathfrak{J} \text{ non-ram.} \\ N\mathfrak{J} \leq x}} \varepsilon_C\left(\left[\frac{\mathbb{L}/\mathbb{K}}{\mathfrak{J}}\right]\right) \Lambda_{\mathbb{K}}(\mathfrak{J}). \end{aligned}$$

In [5] we have proved the following explicit bound.

Theorem. *Assume GRH holds. Let $x \geq 1$, then*

$$(1.1) \quad \left| \frac{|G|}{|C|} \psi_C(x) - x \right| \leq \sqrt{x} \left[\left(\frac{\log x}{2\pi} + 2 \right) \log \Delta_{\mathbb{L}} + \left(\frac{\log^2 x}{8\pi} + 2 \right) n_{\mathbb{L}} \right].$$

Date: May 10, 2023. File name: `pdfupper-k.tex`.

2010 Mathematics Subject Classification. Primary 11R42, Secondary 11Y70.

This result concludes a quite long set of similar but partial computations, originated with Jeffrey Lagarias and Andrew Odlyzko's paper [7] where this result is proved with undetermined constants, and which was followed by the result announced by Joseph Oesterlé [12] and the one of Bruno Winckler [15, Th. 8.1] (both with the same generality and explicit but larger constants), the one of Lowell Schoenfeld [14] (same bound but only for the case $\mathbb{L} = \mathbb{K} = \mathbb{Q}$), and our recent paper [4] (same conclusion, but only for the case $\mathbb{L} = \mathbb{K}$).

Bound (1.1) implies that for every class C there is a prime ideal \mathfrak{p} with

$$(1.2) \quad N\mathfrak{p} \leq \left(\left(\frac{1}{2\pi \log \delta} + o(1) \right) \log \Delta_{\mathbb{L}} (\log \log \Delta_{\mathbb{L}})^2 \right)^2$$

which is not ramified and for which $\left[\frac{\mathbb{L}/\mathbb{K}}{\mathfrak{p}} \right] = C$, where δ is any lower bound for the root discriminant of the family of fields for which we are interested to apply the result: $\sqrt{3}$ is a possible value for all fields, and $\sqrt[3]{23}$ is another possible value when the six quadratic fields $\mathbb{Q}[\sqrt{d}]$ with $d \in \{-7, -3, -2, -1, 2, 5\}$ are excluded.

This consequence of any bound similar to (1.1) is already discussed in Lagarias and Odlyzko's paper, where in fact the existence of a bound of the form $c(\log \Delta_{\mathbb{L}} (\log \log \Delta_{\mathbb{L}})^2)^2$ for some computable (but not explicit) constant c is proved.

The appearance of the factor $(\log \log \Delta_{\mathbb{L}})^4$ in (1.2) is a consequence of the use of (1.1), which actually is not designed for that purpose. In fact, in essence, this comes down to the fact that the kernel $\frac{x^s}{s}$, needed to relate ψ_C to a convenient sum of logarithmic derivatives of Artin L -functions, does not decay very quickly along the vertical lines. To overcome this problem, the authors of [7] also sketched a different approach replacing $\frac{x^s}{s}$ with the kernel $\left(\frac{y^{s-1} - x^{s-1}}{s-1} \right)^2$. With a suitable choice of the parameters x and y in terms of $\log \Delta_{\mathbb{L}}$, this kernel allows to remove the factor $(\log \log \Delta_{\mathbb{L}})^4$ from the bound.

This improvement is not exclusive of this specific kernel, and the same conclusion may be achieved also via different kernels, provided that they decay quickly enough along vertical lines. In particular, we have obtained (1.1) as a by-product of computations for $\psi_C^{(1)}(x) := \sum_{n \leq x} \psi_C(n)$, which is related to the kernel $\frac{x^{s+1}}{s(s+1)}$. Its decay along vertical lines is better than the one of $\frac{x^s}{s}$ and is actually strong enough to get a bound of the type of (1.2), without the factor $(\log \log \Delta_{\mathbb{L}})^4$. In fact, we prove here the following claim as a consequence of some of the inner results we got in [5].

Theorem. *Assume GRH holds. Fix any class C and any integer $k \geq 0$. Assume*

$$\sqrt{x} \geq 1.075 \log \Delta_{\mathbb{L}} + \sqrt{2 \frac{|G|}{|C|} k \log \left(\frac{|G|}{|C|} k \right) + 2 \frac{|G|}{|C|} + 15},$$

where $k \log k$ is set to 0 for $k = 0$. Then $\pi_C(x) \geq k+1$. In other words, if we order the prime ideals $\{\mathfrak{p}_k\}_{k=1}^{\infty}$ which are not ramified and for which $\left[\frac{\mathbb{L}/\mathbb{K}}{\mathfrak{p}} \right] = C$ according to their norm, for every $k \geq 0$ we have

$$N\mathfrak{p}_{k+1} \leq \left(1.075 \log \Delta_{\mathbb{L}} + \sqrt{2 \frac{|G|}{|C|} k \log \left(\frac{|G|}{|C|} k \right) + 2 \frac{|G|}{|C|} + 15} \right)^2.$$

Thus, for instance, there is a non-ramified prime ideal \mathfrak{p} in C with $N\mathfrak{p} \leq \left(1.075 \log \Delta_{\mathbb{L}} + 2 \frac{|G|}{|C|} + 15 \right)^2$ (case $k = 0$) and two such ideals within $\left(1.075 \log \Delta_{\mathbb{L}} + \sqrt{2 \frac{|G|}{|C|} \log \left(\frac{|G|}{|C|} \right) + 2 \frac{|G|}{|C|} + 15} \right)^2$ (case $k = 1$).

The proof of this theorem shows that the constant 15 can be removed when the degree of the field is large enough, but the main constant 1.075 is rooted in the method and can be improved only marginally. In particular it remains larger than 1. This implies that the case $k = 0$ of the theorem is weaker than the analogous conclusion of the paper by Eric Bach and Jonathan Sorenson [1, Th. 3.1], further improved for the case where $\mathbb{K} = \mathbb{Q}$ and \mathbb{L}/\mathbb{Q} is abelian by Youness Lamzouri, Xiannan Li and Kannan Soundararajan [8, Th. 1.2] (see also [9]).

The claim giving more ideals (i.e., $k \geq 1$) cannot be reached with Lagarias–Odlyzko’s, Bach–Sorenson’s or Lamzouri–Li–Soundararajan’s approaches.

The case where $\mathbb{K} = \mathbb{Q}$ and C is the trivial class has been considered also in [3, Corollary 2.1], with similar conclusions, in particular with the same constant for $\log \Delta_{\mathbb{L}}$ but a larger one for the $k \log k$ term.

Finally, we notice that if the field extension \mathbb{L}/\mathbb{K} and the class C are fixed and only the dependence on k is retained, then the theorem says that the norm of the k th prime ideals in C is $\leq (2+o(1))\frac{|G|}{|C|}k \log k$: this is the correct upper bound in its dependency on k and on the density factor $\frac{|G|}{|C|}$, but we know from the prime ideals density theorems that the absolute constant 2 could be 1. This overestimation represents the price we pay in order to get a uniform and totally explicit result.

Acknowledgements. The authors are members of the INdAM group GNSAGA.

2. PRELIMINARY FACTS

For any prime ideal $\mathfrak{p} \subseteq \mathcal{O}_{\mathbb{K}}$, possibly ramified, let \mathfrak{P} be any prime ideal dividing $\mathfrak{p}\mathcal{O}_{\mathbb{L}}$, let I be the inertia group of \mathfrak{P} and τ be one of the Frobenius automorphisms corresponding to $\mathfrak{P}/\mathfrak{p}$. Let

$$\theta(C; \mathfrak{p}^m) := \frac{1}{|I|} \sum_{a \in I} \varepsilon_C(\tau^m a).$$

Notice that $\theta(C; \mathfrak{p}^m) \in [0, 1]$, and that $\theta(C; \mathfrak{p}^m) = \varepsilon_C\left(\left[\frac{\mathbb{L}/\mathbb{K}}{\mathfrak{p}^m}\right]\right)$ for any non-ramified prime \mathfrak{p} and power m . Thus $\theta(C; \cdot)$ extends $\varepsilon_C\left(\left[\frac{\mathbb{L}/\mathbb{K}}{\cdot}\right]\right)$ to ramifying prime ideals powers. With $\theta(C; \cdot)$ at our disposal we define the new function

$$\psi(C; x) := \sum_{\substack{\mathfrak{J} \subseteq \mathcal{O}_{\mathbb{K}} \\ N\mathfrak{J} \leq x}} \theta(C; \mathfrak{J}) \Lambda_{\mathbb{K}}(\mathfrak{J}).$$

Observe that $\psi_C(x)$ and $\psi(C; x)$ are essentially equivalent since they agree except on ramified-prime-powers ideals. However, $\psi(C; \cdot)$ is easier to deal with, since $\theta(C; \cdot)$ is well defined for every prime ideal.

We further set

$$\psi^{(1)}(C; x) := \int_0^x \psi(C; t) dt$$

and, for $s > 1$,

$$K(C; s) := \sum_{\mathfrak{J} \subseteq \mathcal{O}_{\mathbb{K}}} \theta(C; \mathfrak{J}) \Lambda_{\mathbb{K}}(\mathfrak{J}) (N\mathfrak{J})^{-s}.$$

As in [6, Ch. IV Sec. 4, p. 73] and [7, Sec. 5], we have the integral representation

$$(2.1) \quad \psi^{(1)}(C; x) = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} K(C; s) \frac{x^{s+1}}{s(s+1)} ds.$$

The function $\theta(C; \cdot)$ is a class function and therefore can be written as a linear combination of characters of irreducible representations of the group G . A clever trick (due to Deuring [2] and MacCluer [10], see also Lagarias and Odlyzko [7, Lemma 4.1] and [5, p. 445–446]) allows to write this function as a linear combination of characters which are induced from characters of a certain cyclic subgroup H of G specified below. Namely,

$$(2.2) \quad K(C; s) = -\frac{|C|}{|G|} \sum_{\chi} \bar{\chi}(g) \frac{L'}{L}(s, \chi, \mathbb{L}/\mathbb{E}),$$

where g is any fixed element in C , $\mathbb{E} := \mathbb{L}^H$ is the subfield of \mathbb{L} fixed by $H := \langle g \rangle$, $L(s, \chi, \mathbb{L}/\mathbb{E})$ is the Artin L -function associated with the extension \mathbb{L}/\mathbb{E} and the character χ , and the sum runs on all irreducible characters χ of H . Since the extension is abelian, this coincides with a suitable Hecke L -function, by class field theory.

With (2.1), (2.2) produces the identity

$$(2.3) \quad \frac{|G|}{|C|} \psi^{(1)}(C; x) = -\sum_{\chi} \bar{\chi}(g) \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \frac{L'}{L}(s, \chi, \mathbb{L}/\mathbb{E}) \frac{x^{s+1}}{s(s+1)} ds.$$

Finally, we introduce a special notation for the type of sum on characters as the one appearing in (2.3), and for any $f: \widehat{\text{Gal}(\mathbb{L}/\mathbb{E})} \rightarrow \mathbb{C}$ we set

$$\mathcal{M}_C f := \sum_{\chi} \bar{\chi}(g) f(\chi).$$

With this language, Equality (2.3) reads

$$(2.4) \quad \frac{|G|}{|C|} \psi^{(1)}(C; x) = \mathcal{M}_C I_{\chi}(x),$$

where

$$(2.5) \quad I_{\chi}(x) := -\frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \frac{L'}{L}(s, \chi, \mathbb{L}/\mathbb{E}) \frac{x^{s+1}}{s(s+1)} ds.$$

3. SOME COMPUTATIONS WITH ABELIAN ARTIN L -FUNCTIONS

Let $\mathbb{E} \subseteq \mathbb{L}$ be an abelian extension of fields and let χ be any irreducible character of $\text{Gal}(\mathbb{L}/\mathbb{E})$. We will use $L(s, \chi)$ to denote $L(s, \chi, \mathbb{L}/\mathbb{E})$. Also, set $\delta_{\chi} = 1$ if χ is the trivial character, and 0 otherwise.

We recall that for each χ there exist non-negative integers a_{χ}, b_{χ} such that

$$a_{\chi} + b_{\chi} = n_{\mathbb{E}}$$

and a positive integer $Q(\chi)$ such that if we define

$$(3.1) \quad \Gamma_{\chi}(s) := \left[\pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) \right]^{a_{\chi}} \left[\pi^{-\frac{s+1}{2}} \Gamma\left(\frac{s+1}{2}\right) \right]^{b_{\chi}}$$

and

$$(3.2) \quad \xi(s, \chi) := [s(s-1)]^{\delta_{\chi}} Q(\chi)^{s/2} \Gamma_{\chi}(s) L(s, \chi),$$

then $\xi(s, \chi)$ satisfies the functional equation

$$(3.3) \quad \xi(1-s, \bar{\chi}) = W(\chi)\xi(s, \chi),$$

where $W(\chi)$ is a certain constant of absolute value 1. Furthermore, $\xi(s, \chi)$ is an entire function (by class field theory) of order 1 and does not vanish at $s = 0$, and hence by Hadamard's product theorem we have

$$(3.4) \quad \xi(s, \chi) = e^{A_\chi + B_\chi s} \prod_{\rho \in Z_\chi} \left(1 - \frac{s}{\rho}\right) e^{s/\rho}$$

for some constants $A(\chi)$ and $B(\chi)$, where Z_χ is the set of zeros (multiplicity included) of $\xi(s, \chi)$. They are precisely those zeros $\rho = \beta + i\gamma$ of $L(s, \chi)$ for which $0 < \beta < 1$, the so-called “non-trivial zeros” of $L(s, \chi)$. From now on ρ will denote a non-trivial zero of $L(s, \chi)$.

Differentiating (3.2) and (3.4) logarithmically we obtain the identity

$$(3.5) \quad \frac{L'}{L}(s, \chi) = B_\chi + \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho}\right) - \frac{1}{2} \log Q(\chi) - \delta_\chi \left(\frac{1}{s} + \frac{1}{s-1}\right) - \frac{\Gamma'_\chi}{\Gamma_\chi}(s),$$

valid identically in the complex variable s .

Using (3.2), (3.3) and (3.5) one sees that

$$(3.6) \quad \begin{aligned} \frac{L'}{L}(s, \chi) &= \frac{a_\chi - \delta_\chi}{s} + r_\chi + O(s) \quad \text{as } s \rightarrow 0, \\ \frac{L'}{L}(s, \chi) &= \frac{b_\chi}{s+1} + r'_\chi + O(s+1) \quad \text{as } s \rightarrow -1, \end{aligned}$$

where

$$(3.7) \quad r_\chi = B_\chi + \delta_\chi - \frac{1}{2} \log \frac{Q(\chi)}{\pi^{n_\mathbb{E}}} - \frac{a_\chi}{2} \frac{\Gamma'}{\Gamma}(1) - \frac{b_\chi}{2} \frac{\Gamma'}{\Gamma}\left(\frac{1}{2}\right),$$

$$(3.8) \quad r'_\chi = -\frac{L'}{L}(2, \bar{\chi}) - \log \frac{Q(\chi)}{\pi^{n_\mathbb{E}}} - \frac{n_\mathbb{E}}{2} \frac{\Gamma'}{\Gamma}\left(\frac{3}{2}\right) - \frac{n_\mathbb{E}}{2} \frac{\Gamma'}{\Gamma}(1).$$

Comparing (3.7) and (3.5) with $s = 2$, we further get

$$(3.9) \quad r_\chi = \frac{L'}{L}(2, \chi) - \sum_{\rho} \frac{2}{\rho(2-\rho)} + \frac{5}{2} \delta_\chi + b_\chi.$$

Shifting the axis of integration in (2.5) arbitrarily far to the left, we collect the terms coming from the pole of L at $s = 1$ (if any), the non-trivial zeros, the pole of the kernel (and of L'/L , if any) at $s = 0$, the pole of the kernel (and of L'/L , if any) at $s = -1$ and all the remaining terms coming from the trivial zeros of L . This procedure gives the identity

$$(3.10) \quad I_\chi(x) = \delta_\chi \frac{x^2}{2} - \sum_{\rho \in Z_\chi} \frac{x^{\rho+1}}{\rho(\rho+1)} - x r_\chi + r'_\chi + R_\chi(x) \quad \forall x > 1,$$

where r_χ and r'_χ are defined in (3.6) and $R_\chi(x)$ is the explicit function

$$\begin{aligned} f_1(x) &:= \sum_{r=1}^{\infty} \frac{x^{1-2r}}{2r(2r-1)}, & f_2(x) &:= \sum_{r=2}^{\infty} \frac{x^{2-2r}}{(2r-1)(2r-2)}, \\ R_\chi(x) &:= -(a_\chi - \delta_\chi)(x \log x - x) + b_\chi(\log x + 1) - a_\chi f_1(x) - b_\chi f_2(x). \end{aligned}$$

(with $x > 1$). The correctness of this procedure is proved in a way similar to [7, § 6], further simplified by the fact that the integral is absolutely convergent on vertical lines (see also [6, Ch. IV Sec. 4, p. 73]).

According to (2.4), in order to proceed we need to know the effect of the \mathcal{M}_C operator on each term in (3.10). To this effect, we recall a few lemmas that we will need in the following.

Lemma 3.1 ([5, Lemma 1]). *Let*

$$\mathbf{S} := \begin{cases} r_1(\mathbb{L}) + r_2(\mathbb{L}) & \text{if } g \text{ has order 1,} \\ r_2(\mathbb{L}) - 2r_2(\mathbb{E}) & \text{if } g \text{ has order 2,} \\ 0 & \text{otherwise.} \end{cases}$$

Moreover let δ_C be defined to be 1 if C is the trivial class and 0 otherwise. Then

$$\begin{aligned} \mathcal{M}_C a_\chi &= \sum_x \bar{\chi}(g) a_\chi = \mathbf{S}, \\ \mathcal{M}_C b_\chi &= \sum_x \bar{\chi}(g) b_\chi = \delta_C n_{\mathbb{E}} - \mathbf{S} = \delta_C n_{\mathbb{L}} - \mathbf{S}. \end{aligned}$$

From now on, we assume that \mathbb{L}/\mathbb{E} is cyclic, and let Z be the multiset of zeros of the Dedekind zeta function $\zeta_{\mathbb{L}}$. Thus Z is the disjoint union of the sets Z_χ for $\chi \in \widehat{\text{Gal}(\mathbb{L}/\mathbb{E})}$.

Lemma 3.2 ([5, Lemma 2]). *Let f be any complex function with $\sum_{\rho \in Z} |f(\rho)| < \infty$. Then*

$$\mathcal{M}_C \sum_{\rho \in Z_\chi} f(\rho) = \sum_{\rho \in Z} \epsilon(\rho) f(\rho)$$

where, for any $\rho \in Z$, $|\epsilon(\rho)| = 1$ and $\epsilon(\bar{\rho}) = \overline{\epsilon(\rho)}$.

The following lemma comes from (3.9) and Lemmas 3.1 and 3.2.

Lemma 3.3 ([5, Lemma 3]).

$$\mathcal{M}_C r_\chi = 2 \sum_{\rho \in Z} \frac{\epsilon(\rho)}{\rho(2-\rho)} - \frac{n_{\mathbb{L}}}{n_{\mathbb{K}}|C|} \sum_{\mathfrak{J} \subseteq \mathcal{O}_{\mathbb{K}}} \theta(C; \mathfrak{J}) \frac{\Lambda_{\mathbb{K}}(\mathfrak{J})}{(N\mathfrak{J})^2} + n_{\mathbb{L}} \delta_C - \mathbf{S} + \frac{5}{2}.$$

Lemma 3.4 ([5, Lemma 5]). *Define for any $x > 1$, $R_C(x) := \mathcal{M}_C R_\chi(x)$. Then*

$$R_C(x) = \int_0^x \log u \, du - \mathbf{S} \int_1^{x+1} \log u \, du + \delta_C \frac{n_{\mathbb{L}}}{2} \left[\log(x^2 - 1) + x \log \left(\frac{x+1}{x-1} \right) \right].$$

Lemma 3.5 ([5, Lemma 10]). *Assume GRH. Then*

$$\sum_{\rho \in Z} \frac{1}{|\rho(\rho+1)|} \leq 0.5375 \log \Delta_{\mathbb{L}} - 1.0355 n_{\mathbb{L}} + 5.3879 - 0.2635 r_1(\mathbb{L}).$$

We finally prove three technical lemmas.

Lemma 3.6. *Assume GRH. Then*

$$\mathcal{M}_C r_\chi \leq 1.075 \log \Delta_{\mathbb{L}} - 1.571 n_{\mathbb{L}} + 13.276.$$

Proof. By Lemma 3.3, we have

$$\mathcal{M}_C r_\chi \leq 2 \sum_{\rho \in Z} \frac{1}{|\rho(2-\rho)|} + n_{\mathbb{L}} \delta_C - \mathbf{S} + \frac{5}{2}.$$

A brief check shows that $n_{\mathbb{L}} \delta_C - \mathbf{S} \leq r_2(\mathbb{L}) \leq \frac{1}{2} n_{\mathbb{L}}$. Moreover, $|\rho(2-\rho)| = |\rho(\rho+1)|$, thus Lemma 3.5 applies here and the result follows. \square

Lemma 3.7. *We have*

$$-\mathcal{M}_C r'_\chi \leq \log \Delta_{\mathbb{L}}.$$

Proof. As a consequence of (3.8), we have

$$-\mathcal{M}_C r'_\chi = \mathcal{M}_C \frac{L'}{L}(2, \bar{\chi}) + \mathcal{M}_C \log Q(\chi) - n_{\mathbb{E}} \left(\log \pi - \frac{1}{2} \frac{\Gamma'}{\Gamma} \left(\frac{3}{2} \right) - \frac{1}{2} \frac{\Gamma'}{\Gamma}(1) \right) \mathcal{M}_C 1.$$

Letting C_1 to be the class of g^{-1} , we see from (2.2) that

$$\mathcal{M}_C \frac{L'}{L}(2, \bar{\chi}) = -\frac{|G|}{|C|} \overline{K(C_1; 2)}$$

which, by definition of K , is a negative real. Moreover,

$$|\mathcal{M}_C \log Q(\chi)| = \left| \sum_{\chi} \bar{\chi}(g) \log Q(\chi) \right| \leq \sum_{\chi} \log Q(\chi) = \log \Delta_{\mathbb{L}},$$

by the product formula for conductors. The result follows because $n_{\mathbb{E}} \mathcal{M}_C 1 = n_{\mathbb{L}} \delta_C \geq 0$ and $\log \pi - \frac{1}{2} \frac{\Gamma'}{\Gamma} \left(\frac{3}{2} \right) - \frac{1}{2} \frac{\Gamma'}{\Gamma}(1) = 1.41 \dots$ is positive. \square

Lemma 3.8. *If $\mathbb{L} \neq \mathbb{Q}$, for any $x > 1$,*

$$-R_C(x) \leq (n_{\mathbb{L}} - 1) \int_1^{x+1} \log u \, du.$$

Proof. Consider the formula for $R_C(x)$ given in Lemma 3.4. When $r_2(\mathbb{L}) \geq 1$ we have $\mathbf{S} \leq r_1(\mathbb{L}) + r_2(\mathbb{L}) = n_{\mathbb{L}} - r_2(\mathbb{L}) \leq n_{\mathbb{L}} - 1$ producing

$$-R_C(x) \leq \mathbf{S} \int_1^{x+1} \log u \, du \leq (n_{\mathbb{L}} - 1) \int_1^{x+1} \log u \, du.$$

On the other hand, if $r_2(\mathbb{L}) = 0$ then when $\delta_C = 0$ we have $\mathbf{S} = 0$ and $R_C(x) > 0$, while if $\delta_C = 1$ we have $\mathbf{S} = r_1(\mathbb{L}) = n_{\mathbb{L}}$, $\frac{n_{\mathbb{L}}}{2} \delta_C \geq 1$ because $\mathbb{L} \neq \mathbb{Q}$, and

$$\begin{aligned} -R_C(x) &\leq n_{\mathbb{L}} \int_1^{x+1} \log u \, du - \int_0^x \log u \, du - \left[\log(x^2 - 1) + x \log \left(\frac{x+1}{x-1} \right) \right] \\ &= (n_{\mathbb{L}} - 1) \int_1^{x+1} \log u \, du - \log(x-1) - x \log \left(\frac{x}{x-1} \right) \leq (n_{\mathbb{L}} - 1) \int_1^{x+1} \log u \, du. \quad \square \end{aligned}$$

4. PROOF OF THE THEOREM

When $\mathbb{L} = \mathbb{Q}$ the claim follows easily by Chebyshev's bound $\pi(x) \geq \frac{x}{2 \log x}$. For the next computations we assume $\mathbb{L} \neq \mathbb{Q}$.

Lemma 4.1. *Let $x \geq 400$ and $y > 0$, then*

$$(x-y) \log y \leq x(\log x - \log(2 \log x)).$$

Proof. Let $f_x(y) := (x-y) \log y$. Its maximum is attained at a unique point $y_0(x) \in (1, x)$, with $y_0(\log y_0 + 1) = x$. The formula shows that y_0 grows as a function of x . A simple computation shows that

$$\frac{f_x(y_0)}{x} - \log x + \log \log x = g(1 + \log y_0),$$

where

$$g(z) := \log \left(1 + \frac{\log z - 1}{z} \right) + \frac{1}{z} - 1.$$

This function decreases for $z \geq e$ and is lower than $-\log 2$ when $z \geq 5.3193$. Since $5.3193e^{4.3193} = 399.67\dots$, the claim is proved. \square

Let $a_C(n) := \#\{\mathfrak{p} : \mathfrak{p} \text{ unramified, } [\mathbb{L}/\mathbb{K}]_{\mathfrak{p}} = C, N\mathfrak{p} = n\}$ and let also

$$\vartheta_C(x) := \sum_{\substack{\mathfrak{p} \text{ non-ram.} \\ N\mathfrak{p} \leq x}} \varepsilon_C \left(\left[\frac{\mathbb{L}/\mathbb{K}}{\mathfrak{p}} \right] \right) \log N\mathfrak{p} = \sum_{n \leq x} a_C(n) \log n, \quad \vartheta_C^{(1)}(x) := \int_0^x \vartheta_C(t) dt.$$

Then, by Lemma 4.1, for $x \geq 400$,

$$\begin{aligned} \vartheta_C^{(1)}(x) &= \sum_{n \leq x} a_C(n) (x-n) \log n \leq x(\log x - \log(2 \log x)) \sum_{n \leq x} a_C(n) \\ (4.1) \quad &= \pi_C(x) x (\log x - \log(2 \log x)). \end{aligned}$$

Now we produce a lower bound for $\vartheta_C^{(1)}(x)$ out of a lower bound for $\psi^{(1)}(C; x)$.

To ease the notation we set $g_c := |G|/|C|$ and observe that this is a positive integer.

By (2.4), (3.10) and Lemma 3.2, we get

$$g_c \psi^{(1)}(C; x) = \mathcal{M}_C I_\chi(x) = \frac{x^2}{2} - \sum_{\rho \in Z} \epsilon(\rho) \frac{x^{\rho+1}}{\rho(\rho+1)} - x \mathcal{M}_C r_\chi + \mathcal{M}_C r'_\chi + R_C(x)$$

which with the GRH assumption yields

$$\frac{x^2}{2} - g_c \psi^{(1)}(C; x) \leq x^{3/2} \sum_{\rho \in Z} \frac{1}{|\rho(\rho+1)|} + x \mathcal{M}_C r_\chi - \mathcal{M}_C r'_\chi - R_C(x).$$

With Lemmas 3.5–3.8, this gives

$$\begin{aligned} \frac{x^2}{2} - g_c \psi^{(1)}(C; x) &\leq (0.5375(x^{3/2} + 2x) + 1) \log \Delta_{\mathbb{L}} + x^{3/2} (5.4 - 1.0355 n_{\mathbb{L}}) \\ &\quad + \left(\int_1^{x+1} \log u \, du - 1.571x \right) n_{\mathbb{L}} + 13.276x - \int_1^{x+1} \log u \, du. \end{aligned}$$

When $x \geq 400$ the term in $n_{\mathbb{L}}$ appearing in the last line is bounded by $n_{\mathbb{L}} x (\log x - 2.55)$ and the sum of the last two terms by $8.3x$. We further simplify their contribution noticing that

$$n_{\mathbb{L}} x (\log x - 2.55) + 8.3x = n_{\mathbb{L}} x (\log x - 2.4) - 0.15 n_{\mathbb{L}} x + 8.3x \leq n_{\mathbb{L}} x \log x - 2.4 n_{\mathbb{L}} x + 8x,$$

where in the last step we used that $n_{\mathbb{L}} \geq 2$. We thus have

$$\begin{aligned} \frac{x^2}{2} - g_c \psi^{(1)}(C; x) &\leq (0.5375(x^{3/2} + 2x) + 1) \log \Delta_{\mathbb{L}} + x^{3/2} (5.4 - 1.0355 n_{\mathbb{L}}) \\ &\quad + n_{\mathbb{L}} x \log x - 2.4 n_{\mathbb{L}} x + 8x. \end{aligned}$$

Now we remove the contribution to $\psi^{(1)}(C; x)$ of the prime powers \mathfrak{p}^m with $m \geq 2$. Let

$$\vartheta(C; x) := \sum_{\substack{\mathfrak{p} \subset \mathcal{O}_{\mathbb{K}} \\ N\mathfrak{p} \leq x}} \theta(C; \mathfrak{p}) \log(N\mathfrak{p}), \quad \vartheta^{(1)}(C; x) := \int_0^x \vartheta(C; t) dt.$$

The estimation in [13, Th. 13] gives $0 \leq \psi^{(1)}(C; x) - \vartheta^{(1)}(C; x) \leq 1.43\frac{2}{3}x^{3/2}n_{\mathbb{K}}$. Thus

$$\begin{aligned} \frac{x^2}{2} - g_c \vartheta^{(1)}(C; x) &\leq (0.5375x^{3/2} + 1.075x + 1) \log \Delta_{\mathbb{L}} \\ &\quad + x^{3/2}(5.4 - 0.082n_{\mathbb{L}}) + n_{\mathbb{L}}x \log x - 2.4n_{\mathbb{L}}x + 8x \end{aligned}$$

which simplifies to

$$(4.2) \quad \frac{x^2}{2} - g_c \vartheta^{(1)}(C; x) \leq (0.5375x^{3/2} + 1.075x + 1) \log \Delta_{\mathbb{L}} + 2n_{\mathbb{L}}x + 5.4x^{3/2} + 8x,$$

because $(-0.082x^{3/2} + x \log x - 2.4x)/x$ has a maximum at $x = (2/0.082)^2 = 594.88\dots$ where it is lower than 2. The quantities $\vartheta(C; x)$ and $\vartheta_C(x)$ differ only by the contribution of the ramified prime ideals to $\vartheta(C; x)$. In fact,

$$0 \leq \vartheta(C; x) - \vartheta_C(x) \leq \sum_{\substack{\mathfrak{p} \\ \mathfrak{p} \text{ ram.} \\ N\mathfrak{p} \leq x}} \log N\mathfrak{p} \leq \sum_{\mathfrak{p} \text{ ram.}} \log N\mathfrak{p} \leq \log(N\Delta_{\mathbb{L}/\mathbb{K}}) \leq \log \Delta_{\mathbb{L}}.$$

Hence,

$$0 \leq \vartheta^{(1)}(C; x) - \vartheta_C^{(1)}(x) \leq (x-1) \log \Delta_{\mathbb{L}},$$

which with (4.2) gives

$$(4.3) \quad \frac{x^2}{2} - g_c \vartheta_C^{(1)}(x) \leq (0.5375x^{3/2} + g_c x + 1.075x) \log \Delta_{\mathbb{L}} + 2n_{\mathbb{L}}x + 5.4x^{3/2} + 8x.$$

By (4.1) and (4.3), in order to have $\pi_C(x) > k$ it is sufficient to have

$$\frac{x^2}{2} > (0.5375x^{3/2} + g_c x + 1.075x) \log \Delta_{\mathbb{L}} + 2n_{\mathbb{L}}x + 5.4x^{3/2} + 8x + k g_c x (\log x - \log(2 \log x)),$$

i.e.

$$(4.4) \quad \sqrt{x} > \left(1.075 + \frac{2g_c + 2.15}{\sqrt{x}}\right) \log \Delta_{\mathbb{L}} + 4 \frac{n_{\mathbb{L}}}{\sqrt{x}} + 10.8 + \frac{16}{\sqrt{x}} + 2k g_c \frac{\log x - \log(2 \log x)}{\sqrt{x}}$$

which is true when

$$\sqrt{x} = 1.075 \log \Delta_{\mathbb{L}} + \sqrt{2g_c k \log(g_c k)} + 2g_c + 15.$$

Proof. Let

$$A := 1.075 \log \Delta_{\mathbb{L}} + 2g_c + 15$$

and

$$B := \left(1.075 + \frac{2g_c + 2.15}{\sqrt{x}}\right) \log \Delta_{\mathbb{L}} + 4 \frac{n_{\mathbb{L}}}{\sqrt{x}} + 10.8 + \frac{16}{\sqrt{x}}.$$

To show that (4.4) holds with the indicated value of x , it is sufficient to prove

$$(4.5) \quad A + \sqrt{2g_c k \log(g_c k)} > B + 2k g_c \frac{\log x - \log(2 \log x)}{\sqrt{x}}.$$

We have

$$(A - B - 1)\sqrt{x} = (2g_c + 3.2)\sqrt{x} - ((2g_c + 2.15) \log \Delta_{\mathbb{L}} + 4n_{\mathbb{L}} + 16) \geq 1.4 \log \Delta_{\mathbb{L}} - 4n_{\mathbb{L}} + 72$$

which is positive, according to entry $b = 4$ in [11, Table 3]. Since $A - B > 1$, our claim will hold if

$$\sqrt{2g_c k \log(g_c k)} + 1 \geq 2g_c k \frac{\log x - \log(2 \log x)}{\sqrt{x}},$$

i.e. $k = 0$ or

$$(4.6) \quad \frac{\sqrt{\log y}}{\sqrt{2y}} + \frac{1}{2y} \geq \frac{\log x - \log(2 \log x)}{\sqrt{x}},$$

where $y := g_c k \geq 1$. The right-hand side decreases if $x \geq 30$ hence is at most 0.2 and the left-hand side is larger than 0.2 for $1 \leq y \leq 120$. We thus assume $y \geq 120$, and in that case $x \geq 2y \log y \geq 30$, hence (4.6) holds if

$$\frac{\sqrt{\log y}}{\sqrt{2y}} \geq \frac{\log(2y \log y) - \log(2 \log(2y \log y))}{\sqrt{2y \log y}}$$

i.e.

$$\log y \geq \log(2y \log y) - \log(2 \log(2y \log y))$$

which is obviously true in this range. \square

This proves the claim under the assumption that $x \geq 400$. The exceptions to this condition are the cases where

$$1.075 \log \Delta_{\mathbb{L}} + \sqrt{2g_c k \log(g_c k)} + 2g_c + 15 < 20$$

and this happens only when $g_c = 1$, $\Delta_{\mathbb{L}} \leq 16$ and $k \leq 2$.

For these remaining cases we check directly the existence of the corresponding ideals. We observe that $g_c = |G|/|C| = 1$ if and only if $|G| = 1$ and hence $\mathbb{L} = \mathbb{K}$. Moreover, $\Delta_{\mathbb{L}} \leq 16$ implies $n_{\mathbb{L}} \leq 2$. Hence it is sufficient to check that, in quadratic fields, there are at least three ideals of norm at most $\lfloor (1.075 \log 3 + 17)^2 \rfloor = 330$. They exist because the primes above 2, 3 and 5 have norm at most 25.

REFERENCES

1. E. Bach and J. Sorenson, *Explicit bounds for primes in residue classes*, Math. Comp. **65** (1996), no. 216, 1717–1735.
2. M. Deuring, *Über den Tschebotareffschen Dichtigkeitssatz*, Math. Ann. **110** (1935), no. 1, 414–415.
3. L. Grenié and G. Molteni, *Explicit smoothed prime ideals theorems under GRH*, Math. Comp. **85** (2016), no. 300, 1875–1899.
4. ———, *Explicit versions of the prime ideal theorem for Dedekind zeta functions under GRH*, Math. Comp. **85** (2016), no. 298, 889–906.
5. ———, *An effective Chebotarev density theorem under GRH*, J. Number Theory **200** (2019), 441–485.
6. A. E. Ingham, *The distribution of prime numbers*, Cambridge University Press, Cambridge, 1990.
7. J. C. Lagarias and A. M. Odlyzko, *Effective versions of the Chebotarev density theorem*, Algebraic number fields: L -functions and Galois properties (Proc. Sympos., Univ. Durham, Durham, 1975), Academic Press, London, 1977, pp. 409–464.
8. Y. Lamzouri, X. Li, and K. Soundararajan, *Conditional bounds for the least quadratic non-residue and related problems*, Math. Comp. **84** (2015), no. 295, 2391–2412.
9. ———, *Corrigendum to “Conditional bounds for the least quadratic non-residue and related problems”*, Math. Comp. **86** (2017), no. 307, 2551–2554.
10. C. R. MacCluer, *A reduction of the Čebotarev density theorem to the cyclic case*, Acta Arith. **15** (1968), 45–47.
11. A. M. Odlyzko, *Discriminant bounds*, <http://www.dtc.umn.edu/~odlyzko/unpublished/index.html>, 1976.
12. J. Oesterlé, *Versions effectives du théorème de Chebotarev sous l’hypothèse de Riemann généralisée*, Astérisque **61** (1979), 165–167.

13. J. B. Rosser and L. Schoenfeld, *Approximate formulas for some functions of prime numbers*, Illinois J. Math. **6** (1962), 64–94.
14. L. Schoenfeld, *Sharper bounds for the Chebyshev functions $\theta(x)$ and $\psi(x)$. II*, Math. Comp. **30** (1976), no. 134, 337–360, *Corrigendum* in Math. Comp. **30** (1976), no. 136, 900.
15. B. Winckler, *Théorème de Chebotarev effectif*, arxiv:1311.5715, <http://arxiv.org/abs/1311.5715>, 2013.

(L. Grenié) DIPARTIMENTO DI INGEGNERIA GESTIONALE, DELL'INFORMAZIONE E DELLA PRODUZIONE, UNIVERSITÀ DI BERGAMO, VIALE MARCONI 5, 24044 DALMINE ITALY

Email address: loic.grenie@gmail.com

(G. Molteni) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI MILANO, VIA SALDINI 50, I-20133 MILANO, ITALY

Email address: giuseppe.molteni1@unimi.it