



UNIVERSITÀ DEGLI STUDI DI BERGAMO
DIPARTIMENTO DI INGEGNERIA DELL'INFORMAZIONE
E METODI MATEMATICI[°]

QUADERNI DEL DIPARTIMENTO

Department of Information Technology and Mathematical Methods

Working Paper

Series “*Mathematics and Statistics*”

n. 9/MS – 2009

F-manifolds and integrable systems

of hydrodynamic type

by

P. Lorenzoni, M. Pedroni, A. Raimondo

COMITATO DI REDAZIONE[§]

Series Information Technology (IT): Stefano Paraboschi
Series Mathematics and Statistics (MS): Luca Brandolini, Ilia Negri

[§] L'accesso alle *Series* è approvato dal Comitato di Redazione. I *Working Papers* della Collana dei Quaderni del Dipartimento di Ingegneria dell'Informazione e Metodi Matematici costituiscono un servizio atto a fornire la tempestiva divulgazione dei risultati dell'attività di ricerca, siano essi in forma provvisoria o definitiva.

F -manifolds and integrable systems of hydrodynamic type

Paolo Lorenzoni*, Marco Pedroni**, Andrea Raimondo***

* Dipartimento di Matematica e Applicazioni
Università di Milano-Bicocca
Via Roberto Cozzi 53, I-20125 Milano, Italy
paolo.lorenzoni@unimib.it

** Dipartimento di Ingegneria dell'informazione e metodi matematici
Università di Bergamo - Sede di Dalmine
viale Marconi 5, I-24044 Dalmine BG, Italy
marco.pedroni@unibg.it

*** Department of Mathematics, Imperial College
180 Queen's Gate, London SW7 2AZ, UK
a.raimondo@imperial.ac.uk

Abstract

We investigate the role of Hertling-Manin condition on the structure constants of an associative commutative algebra in the theory of integrable systems of hydrodynamic type. In such a framework we introduce the notion of F -manifold with compatible connection generalizing a structure introduced by Manin.

1 Introduction

In their seminal papers [8, 25], Dubrovin, Novikov, and Tsarev pointed out a deep relation between the integrability properties of systems of PDEs of hydrodynamic type

$$u_t^i = V_j^i u_x^j, \quad i = 1, \dots, n, \quad (1)$$

(sum over repeated indices is understood) and geometrical—in particular, Riemannian—structures on the target manifold M , where (u^1, \dots, u^n) play the role of coordinates. Probably, the most important of such structures is the notion of Frobenius manifold, introduced by

Dubrovin (see, e.g., [4]) in order to give a coordinate-free description of the famous WDVV equations. A crucial ingredient involved in the definition of Frobenius manifolds is a $(1, 2)$ -type tensor field c giving an associative commutative product on every tangent space:

$$(X \circ Y)^i := c_{jk}^i X^j Y^k ,$$

where X and Y are vector fields. More recently [17], Hertling and Manin showed that this product satisfies the condition

$$\begin{aligned} [X \circ Y, Z \circ W] - [X \circ Y, Z] \circ W - [X \circ Y, W] \circ Z - X \circ [Y, Z \circ W] + X \circ [Y, Z] \circ W + \\ + X \circ [Y, W] \circ Z - Y \circ [X, Z \circ W] + Y \circ [X, Z] \circ W + Y \circ [X, W] \circ Z = 0 , \end{aligned} \quad (2)$$

or, in terms of the components of c ,

$$(\partial_s c_{jl}^k) c_{im}^s + (\partial_j c_{im}^s) c_{sl}^k - (\partial_s c_{im}^k) c_{jl}^s - (\partial_i c_{jl}^s) c_{sm}^k - (\partial_l c_{jm}^s) c_{si}^k - (\partial_m c_{li}^s) c_{js}^k = 0 . \quad (3)$$

They called *F-manifold* a manifold endowed with an associative commutative multiplicative structure satisfying condition (2).

The aim of this paper is to study the properties of the PDEs of hydrodynamic type associated with *F*-manifolds. The system (3) and its relation with integrable systems has been considered from a different point of view in [18]. Here, following the insights coming from the case of the principal hierarchy in the context of Frobenius manifolds, we will assume such PDEs to be of the form

$$u_t^i = (V_X)^i_j u_x^j, \quad i = 1, \dots, n, \quad (V_X)^i_j := c_{jk}^i X^k, \quad (4)$$

where X is a vector field on M and c satisfies (2). These assumptions have two important consequences, spelled out respectively in Section 2 and 3:

1. For any choice of the vector field X , the Haantjes tensor associated with the $(1,1)$ tensor field V_X vanishes.

2. They allow one to write the condition of commutativity of two flows of the form (4) as a simple requirement on the corresponding vector fields on M .

Starting from Section 4, we put into the game an additional structure, namely a connection ∇ satisfying the symmetry condition

$$(\nabla_X c)(Y, Z) = (\nabla_Y c)(X, Z), \quad (5)$$

for all vector fields X , Y , and Z . Remarkably, as shown by Hertling [16], condition (2) follows from (5).

In Section 4, following Manin [20], we study the special case where the connection ∇ is flat and we show how to construct an integrable hierarchy of hydrodynamic type. The construction is divided in two steps. First—using a basis of flat vector fields—one defines a set of flows, known as *primary flows*. Then, from these flows one can define recursively the higher flows of the hierarchy. In this way, each primary flow turns out to be the starting

point of a hierarchy. This construction is a straightforward generalization of the principal hierarchy defined by Dubrovin in the case of Frobenius manifolds [4].

The general (non-flat) case is studied in Section 5, where we introduce the notion of F -manifold with compatible (non-flat) connection ∇ and we show that the associated integrable systems of hydrodynamic type are defined by a family of vector fields satisfying the following condition:

$$c_{jm}^i \nabla_k X^m = c_{km}^i \nabla_j X^m. \quad (6)$$

In the non-flat case the existence of solutions of the above system is not guaranteed. Indeed, we prove that every solution X of (6) satisfies the condition

$$(R_{lmi}^k c_{pk}^n + R_{lip}^k c_{mk}^n + R_{lpm}^k c_{ik}^n) X^l = 0,$$

where R is the curvature tensor of ∇ . It is thus natural to introduce the following requirement on the curvature:

$$R_{lmi}^k c_{pk}^n + R_{lip}^k c_{mk}^n + R_{lpm}^k c_{ik}^n = 0. \quad (7)$$

If the structure constants c_{jk}^i admit canonical coordinates, condition (7) is related to the well-known semi-Hamiltonian property introduced by Tsarev [25] as compatibility condition for the linear system providing the symmetries of a diagonal system of hydrodynamic type.

In Section 6, motivated by the Hamiltonian theory of systems of hydrodynamic type, we consider the case of metric connections and we introduce the notion of Riemannian F -manifold. Finally, in Section 7, we discuss in details an important example: the reductions of the dispersionless KP hierarchy (also known as Benney chain).

2 The Haantjes tensor

An important class of systems of hydrodynamic type, widely studied in the literature, consists in those systems which admit diagonal form. We say that a system (1) is diagonalizable if there exists a set of coordinates (r^1, \dots, r^n) —usually called *Riemann invariants*—such that the tensor V_j^i is diagonal in these coordinates: $V_j^i(r) = v^i \delta_j^i$. Then the system takes the (diagonal) form

$$r_t^i = v^i(r^1, \dots, r^n) r_x^i, \quad i = 1, \dots, n.$$

It is important to recall that there exists an invariant criterion for the diagonalizability. One first introduces the *Nijenhuis tensor* of V as

$$N_V(X, Y) = [VX, VY] - V[X, VY] - V[VX, Y] + V^2[X, Y],$$

where X and Y are arbitrary vector fields, and then defines the *Haantjes tensor* as

$$H_V(X, Y) = N_V(VX, VY) - V N_V(X, VY) - V N_V(VX, Y) + V^2 N_V(X, Y).$$

In the case when V has mutually distinct eigenvalues, then V is diagonalizable if and only if its Haantjes tensor is identically zero. In this section, we consider the Haantjes tensor of

$$(V_Z)_j^i = c_{jk}^i Z^k, \quad (8)$$

where c satisfies the Hertling-Manin condition (2). For a $(1, 1)$ - type tensor field of the form (8), the Nijenhuis tensor reads

$$N_{V_Z}(X, Y) = [Z \circ X, Z \circ Y] + Z^2 \circ [X, Y] - Z \circ [X, Z \circ Y] - Z \circ [Z \circ X, Y].$$

By using the Hertling-Manin condition (2) evaluated at $X = Z$, this can be written as

$$N_{V_Z}(X, Y) = [X \circ Z, Z] \circ Y - [X, Y] \circ Z \circ Y + [Z, Y \circ Z] \circ X - [Z, Y] \circ X \circ Z,$$

using this identity it is easy to prove the following

Theorem 1 *The Haantjes tensor associated with V_Z vanishes for any choice of the vector field Z .*

Proof. Let us write for simplicity N in place of N_{V_Z} . Then, we have that

$$\begin{aligned} H_{V_Z}[X, Y] &= N[Z \circ X, Z \circ Y] + Z^2 \circ N[X, Y] - Z \circ N[X, Z \circ Y] - Z \circ N[Z \circ X, Y] = \\ &= [X \circ Z^2, Z] \circ Y \circ Z - [X \circ Z, Z] \circ Z^2 \circ Y + [Z, Y \circ Z^2] \circ X \circ Z + \\ &- [Z, Y \circ Z] \circ X \circ Z^2 + [X \circ Z] \circ Y \circ Z^2 - [X, Z] \circ Z^3 \circ Y + \\ &+ [Z, Y \circ Z] \circ X \circ Z^2 - [Z, Y] \circ X \circ Z^3 - [X \circ Z, Z] \circ Z^2 \circ Y + \\ &+ [X, Z] \circ Z^3 \circ Y - [Z, Y \circ Z^2] \circ X \circ Z + [Z, Y \circ Z] \circ X \circ Z^2 + \\ &- [X \circ Z^2, Z] \circ Y \circ Z + [X \circ Z, Z] \circ Z^2 \circ Y - [Z, Y \circ Z] \circ X \circ Z^2 + \\ &+ [Z, Y] \circ X \circ Z^3 = 0, \end{aligned}$$

where $Z^2 = Z \circ Z$ and $Z^3 = Z \circ Z \circ Z$. □

Suppose now that X is a vector field such that V_X has everywhere distinct real eigenvalues (v^1, \dots, v^n) . Since the Haantjes tensor of V_X vanishes, there exist local coordinates (r^1, \dots, r^n) such that $(V_X)_j^i = \delta_j^i v^i$. These coordinates are the Riemann invariants of the corresponding system of hydrodynamic type. Moreover, we have

Proposition 2 *The components of the tensor field c in the coordinates (r^1, \dots, r^n) are given by*

$$c_{ij}^k = f_i \delta_i^k \delta_j^k.$$

Furthermore, if $f_j \neq 0$ for all j , then f_i depends on the variable r^i only.

Proof. In diagonal coordinates we have

$$(V_X)_j^i = c_{jk}^i X^k = v^i \delta_j^i,$$

hence, we get

$$c_{pq}^j c_{jk}^i X^k = c_{pq}^j v^i \delta_j^i = c_{pq}^i v^i.$$

On the other hand, due to the associativity of the algebra, we can also write

$$c_{pq}^j c_{jk}^i X^k = c_{pk}^j c_{jq}^i X^k = c_{jq}^i v^j \delta_p^j = c_{pq}^i v^p \quad (\text{no sum over } p),$$

and therefore,

$$c_{pq}^i (v^i - v^p) = 0.$$

Since the algebra is commutative and the eigenvalues of V_X are pairwise distinct, this means that the structure constants, in the coordinates (r^1, \dots, r^n) , take the form

$$c_{jk}^i = f_i \delta_j^i \delta_k^i, \quad (9)$$

where the f_i are arbitrary functions, depending in principle on all the variables r^1, \dots, r^n . The requirement on the structure constants c to satisfy the Hertling-Manin condition (3) implies further constraints on the functions f_i . Indeed, substituting (9) into (3), we get a set of equations the f_i have to satisfy; considering for instance the case $m = j \neq k = i = l$, we get

$$f_j \partial_j f_k = 0,$$

which means that, in the non-degenerate case when $f_j \neq 0$ for all j , then f_k depends on r^k only. It is easy to check that conditions (3) give no further restrictions on the f_i ; the proposition is proved. \square

If the functions f_i are everywhere different from zero, then it is easy to show that there exist local coordinates, called *canonical coordinates*, such that $c_{ij}^k = \delta_i^k \delta_j^k$. Moreover, in this case, the vector field

$$e = \sum_{i=1}^n \frac{1}{f_i} \frac{\partial}{\partial r^i}$$

is globally defined and is the unity of the algebra.

Remark 3 *If the algebra has a unity e , then the Hertling-Manin condition implies*

$$\text{Lie}_e c = 0.$$

Indeed, for $X = Y = e$ the Hertling-Manin condition becomes

$$-[e, Z \circ W] + [e, Z] \circ W + [e, W] \circ Z = 0.$$

Remark 4 *An alternative proof of the existence of canonical coordinates has been given in [17] under the assumption of semisimplicity of the algebra, that is, the existence of a basis of idempotents.*

3 Commutativity of the flows

As a consequence of the Hertling-Manin condition, the conditions for the commutativity of two hydrodynamical flows take a rather simple form.

Proposition 5 *The flows*

$$u_t^i = [V_X]_j^i u_x^j = c_{jk}^i X^j u_x^k \quad (10)$$

and

$$u_\tau^i = [V_Y]_j^i u_x^j = c_{jk}^i Y^j u_x^k \quad (11)$$

commute if and only if the vector fields X and Y satisfy the condition

$$((\text{Lie}_X c)(Y, Z) - (\text{Lie}_Y c)(X, Z) + [X, Y] \circ Z) \circ Z = 0,$$

for any vector field Z . Equivalently,

$$\begin{aligned} & ((\text{Lie}_X c)(Y, Z) - (\text{Lie}_Y c)(X, Z) + [X, Y] \circ Z) \circ W \\ & + ((\text{Lie}_X c)(Y, W) - (\text{Lie}_Y c)(X, W) + [X, Y] \circ W) \circ Z = 0 \end{aligned}$$

for all pairs (Z, W) of vector fields. In local coordinates this means that

$$\begin{aligned} & c_{is}^r \left[(\text{Lie}_X c)_{jq}^i Y^q - (\text{Lie}_Y c)_{jq}^i X^q + c_{jq}^i [X, Y]^q \right] \\ & + c_{ij}^r \left[(\text{Lie}_X c)_{sq}^i Y^q - (\text{Lie}_Y c)_{sq}^i X^q + c_{sq}^i [X, Y]^q \right] = 0. \end{aligned}$$

Proof. It is well-known that the commutativity of the flows (10) and (11) is equivalent to the following requirements:

1. The $(1, 1)$ -tensor fields V_X and V_Y (seen as endomorphism of the tangent bundle) commute.

2. For any vector field Z the following condition is satisfied:

$$[V_X(Z), V_Y(Z)] - V_X([Z, V_Y(Z)]) + V_Y([Z, V_X(Z)]) = 0,$$

that is to say,

$$[Z \circ X, Z \circ Y] - X \circ [Z, Z \circ Y] + Y \circ [Z, Z \circ X] = 0.$$

The first requirement is automatically verified due to the associativity of the algebra. Making use of identity (2), the second one becomes

$$([Z \circ X, Y] + [X, Z \circ Y] - [X, Z] \circ Y - [X, Y] \circ Z - X \circ [Z, Y]) \circ Z = 0. \quad (12)$$

A simple calculation shows that the quantity in the bracket, namely

$$[Z \circ X, Y] + [X, Z \circ Y] - [X, Z] \circ Y - [X, Y] \circ Z - X \circ [Z, Y],$$

is equal to

$$(\text{Lie}_X c)(Y, Z) - (\text{Lie}_Y c)(X, Z) + [X, Y] \circ Z. \quad (13)$$

Substituting (13) into (12), we get the thesis. \square

Corollary 6 *A sufficient condition for the commutativity of the hydrodynamic flows (10) and (11) is that*

$$(\text{Lie}_X c)(Y, Z) - (\text{Lie}_Y c)(X, Z) + [X, Y] \circ Z = 0 \quad (14)$$

for all vector fields Z , that is,

$$(\text{Lie}_X c)_{pq}^i Y^q - (\text{Lie}_Y c)_{pq}^i X^q + c_{pq}^i [X, Y]^q = 0 \quad (15)$$

or, equivalently,

$$\text{Lie}_X V_Y - \text{Lie}_Y V_X - V_{[X, Y]} = 0. \quad (16)$$

4 Dubrovin principal hierarchy

In this section, we adapt Dubrovin's construction of the principal hierarchy [4] to the case of F -manifolds with compatible flat connection introduced by Manin in [20].

Definition 7 *An F -manifold with compatible flat connection is a manifold endowed with an associative commutative multiplicative structure given by a $(1, 2)$ -tensor field c and a flat torsionless connection ∇ satisfying the symmetry condition*

$$\nabla_l c_{jk}^i = \nabla_j c_{lk}^i, \quad (17)$$

meaning that ∇c is totally symmetric:

$$(\nabla_X c)(Y, Z) = (\nabla_Y c)(X, Z), \quad (18)$$

for all vector fields X, Y , and Z .

Notice that Hertling-Manin condition (2) does not appear in the above definition. Indeed, as proved by Hertling in [16], it is a consequence of the existence of a torsionless (even non-flat) connection ∇ satisfying (17).

Remark 8 *Notice that in flat coordinates condition (17) reads*

$$\partial_l c_{jk}^i = \partial_j c_{lk}^i.$$

This, together with the commutativity of the algebra, implies that

$$c_{jk}^i = \partial_j C_k^i = \partial_j \partial_k C^i.$$

Therefore, condition (17) is equivalent to the local existence of a vector field C satisfying, for any pair (X, Y) of flat vector fields, the condition

$$X \circ Y = [X, [Y, C]].$$

The above condition appears in the original definition of Manin [20].

Let us construct now the principal hierarchy. In order to do so, the first step consists in defining the primary flows. Since the connection is flat, we can consider a basis $(X_{(1,0)}, \dots, X_{(n,0)})$ of flat vector fields; the primary flows are thus defined as

$$u_{t_{(p,0)}}^i = c_{jk}^i X_{(p,0)}^k u_x^j. \quad (19)$$

Proposition 9 *The primary flows (19) commute.*

Proof. Since the $X_{(p,0)}$ are flat and the torsion vanishes, they commute and

$$\text{Lie}_{X_{(p,0)}} c = \nabla_{X_{(p,0)}} c.$$

Therefore, the commutativity condition (14) for the vector fields $X = X_{(p,0)}$ and $Y = X_{(q,0)}$ follows from condition (17). \square

Starting from the primary flows (19) one can introduce the “higher flows” of the hierarchy, defined as

$$u_{t_{(p,\alpha)}}^i = c_{jk}^i X_{(p,\alpha)}^j u_x^k, \quad (20)$$

by means of the following recursive relations:

$$\nabla_j X_{(p,\alpha)}^i = c_{jk}^i X_{(p,\alpha-1)}^k. \quad (21)$$

Remark 10 *The flatness of the connection ∇ , the symmetry of the tensor ∇c (condition (17)) and the associativity of the algebra with structure constants c_{jk}^i are equivalent to the flatness of the one-parameter family of connections defined, for any pair of vector fields X and Y , by*

$$\tilde{\nabla}_X Y = \nabla_X Y + z X \circ Y, \quad z \in \mathbb{C}.$$

The vector fields obtained by means of the recursive relations (21) are nothing but the z -coefficients of a basis of flat vector fields of the deformed connection [4].

In order to show that the higher flows (20) are well-defined, it is necessary to prove the following

Proposition 11 *The recursive relations (21) are compatible.*

Proof. We note that the recursive relations (21) can be written in the form

$$\partial_j X_{(p,\alpha)}^i = -\Gamma_{jk}^i X_{(p,\alpha)}^k - c_{kj}^i X_{(p,\alpha-1)}^k,$$

thus, we have

$$\begin{aligned} (\partial_j \partial_m - \partial_m \partial_j) X_{(p,\alpha)}^i &= [\partial_m \Gamma_{jl}^i - \partial_j \Gamma_{ml}^i - \Gamma_{jk}^i \Gamma_{ml}^k + \Gamma_{mk}^i \Gamma_{jl}^k] X_{(p,\alpha)}^l + \\ &[\partial_m c_{jl}^i - \partial_j c_{ml}^i - \Gamma_{kj}^i c_{ml}^k - \Gamma_{lm}^k c_{jk}^i + \Gamma_{km}^i c_{jl}^k + \Gamma_{lj}^k c_{mk}^i] X_{(p,\alpha-1)}^l \\ &+ [c_{jk}^i c_{ml}^k - c_{mk}^i c_{jl}^k] X_{(p,\alpha-2)}^l. \end{aligned}$$

The flatness of the connection ∇ , together with identity (17) and the associativity of the algebra, implies the vanishing of the quantity above. Therefore, relations (21) are compatible. \square

Since the primary flows (19) commute and the recursive relations (21) are compatible, it only remains to prove the following

Theorem 12 *The flows of the principal hierarchy commute.*

Proof. Let us consider the hydrodynamic flows associated with the vector fields $X_{(p,\alpha)}$ and $X_{(q,\beta)}$. In order to show that these flows commute, we prove that they satisfy the sufficient condition (15). In local coordinates it reads:

$$\begin{aligned} & X_{(p,\alpha)}^m (\partial_m c_{jk}^i) X_{(q,\beta)}^k - X_{(q,\beta)}^m (\partial_m c_{jk}^i) X_{(p,\alpha)}^k + \\ & - c_{jk}^l (\partial_l X_{(p,\alpha)}^i) X_{(q,\beta)}^k + c_{lk}^i (\partial_j X_{(p,\alpha)}^l) X_{(q,\beta)}^k + \\ & + c_{jl}^i (\partial_k X_{(p,\alpha)}^l) X_{(q,\beta)}^k + c_{jk}^l (\partial_l X_{(q,\beta)}^i) X_{(p,\alpha)}^k + \\ & - c_{lk}^i (\partial_j X_{(q,\beta)}^l) X_{(p,\alpha)}^k - c_{jl}^i (\partial_k X_{(q,\beta)}^l) X_{(p,\alpha)}^k + \\ & - c_{jk}^i ((\partial_l X_{(p,\alpha)}^k) X_{(q,\beta)}^l + (\partial_l X_{(q,\beta)}^k) X_{(p,\alpha)}^l) = 0. \end{aligned}$$

In particular, if the coordinates are flat, the first row vanishes due to the symmetry of the tensor ∇c . Moreover, using the recursive relations (21) we obtain

$$\begin{aligned} & -c_{jk}^l c_{ln}^i X_{(p,\alpha-1)}^n X_{(q,\beta)}^k + c_{lk}^i c_{jn}^l X_{(p,\alpha-1)}^n X_{(q,\beta)}^k + \\ & + c_{jl}^i c_{kn}^l X_{(p,\alpha-1)}^n X_{(q,\beta)}^k + c_{jk}^l c_{ln}^i X_{(q,\beta-1)}^n X_{(p,\alpha)}^k + \\ & - c_{lk}^i c_{jn}^l X_{(q,\beta-1)}^n X_{(p,\alpha)}^k - c_{jl}^i c_{kn}^l X_{(q,\beta-1)}^n X_{(p,\alpha)}^k + \\ & - c_{jk}^i c_{mn}^k X_{(p,\alpha-1)}^n X_{(q,\beta)}^m + c_{jk}^i c_{mn}^k X_{(q,\beta-1)}^n X_{(p,\alpha)}^m \end{aligned}$$

which vanishes due to the associativity of the algebra. \square

Remark 13 *The flows of the principal hierarchy are well-defined even in the case when the torsion of ∇ does not vanish. However, their commutativity depends crucially on this additional assumption.*

5 F -manifolds with compatible connection and related integrable systems

From the point of view of the theory of integrable systems of hydrodynamic type, the “flat case” and the associated principal hierarchy are exceptional. Therefore, it is quite natural to extend the notion of F -manifolds with compatible flat connection to the non-flat case. As a starting point, we consider an F -manifold endowed with a connection ∇ satisfying (17). If ∇ is flat, we know how to construct integrable systems of hydrodynamic type. Indeed, the

starting point of the construction of the previous section is a basis of *flat* vector fields, and the recursive procedure (21) defining the “higher” vector fields and the corresponding flows is well-defined as a consequence of the vanishing of the curvature. In the non-flat case, in order to define integrable systems of hydrodynamic type one needs to find an alternative way to select the vector fields.

5.1 Hydrodynamic-type systems associated with F -manifolds

In the flat case, the vector fields X defining the principal hierarchy satisfy the condition

$$(\nabla_Z X) \circ W = (\nabla_W X) \circ Z \quad (22)$$

for all pairs (Z, W) of vector fields, that is, in local coordinates,

$$c_{jm}^i \nabla_k X^m = c_{km}^i \nabla_j X^m . \quad (23)$$

Indeed, in the case of the flat vector fields $X_{(p,0)}$ defining the primary flows, both sides of (23) vanish due to

$$\nabla_k X_{(p,0)}^m = 0, \quad p = 1, \dots, n ,$$

while the vector fields defining the higher flows of the hierarchy satisfy (23) due to the associativity of the algebra:

$$c_{jm}^i \nabla_k X_{(p,\alpha)}^m = c_{jm}^i c_{kl}^m X_{(p,\alpha-1)}^l = c_{km}^i c_{jl}^m X_{(p,\alpha-1)}^l = c_{km}^i \nabla_j X_{(p,\alpha)}^m .$$

A crucial remark is the following: if ∇ satisfies condition (17), then *any pair of solutions of (23) defines commuting flows even if the connection ∇ is not flat*. More precisely, we have the following

Proposition 14 *If X and Y are two vector fields satisfying condition (22), then the associated flows*

$$u_t^i = c_{jk}^i X^k u_x^j \quad (24)$$

and

$$u_\tau^i = c_{jk}^i Y^k u_x^j \quad (25)$$

commute.

Proof. Recall from Proposition 5 that the flows (24) and (25) commute if and only if

$$((\text{Lie}_X c)(Y, Z) - (\text{Lie}_Y c)(X, Z) + [X, Y] \circ Z) \circ Z = 0 \quad (26)$$

for any vector field Z . On the other hand, the vanishing of the torsion of ∇ gives the identity

$$(\text{Lie}_X c)(Y, Z) = (\nabla_X c)(Y, Z) - \nabla_{c(Y,Z)} X + c(Y, \nabla_Z X) + c(\nabla_Y X, Z) ,$$

and this, together with the symmetry (18) of ∇c , can be used to write the term in the bracket of (26) as

$$-\nabla_{Y \circ Z} X + \nabla_{X \circ Z} Y + [Y, X] \circ Z.$$

Multiplying the above identity by Z , and using property (22) for the vector fields X and Y , we obtain

$$\begin{aligned} & -(\nabla_{Y \circ Z} X) \circ Z + (\nabla_{X \circ Z} Y) \circ Z + [Y, X] \circ Z^2 = \\ & -(\nabla_Z X) \circ (Y \circ Z) + (\nabla_Z Y) \circ (X \circ Z) + [Y, X] \circ Z^2 = \\ & -(\nabla_Y X) \circ Z^2 + (\nabla_X Y) \circ Z^2 + [Y, X] \circ Z^2 = 0. \end{aligned}$$

The proposition is proved. \square

Remark 15 From (17) and (22) it follows that the $(1,1)$ -tensor field

$$(V_X)_j^i = c_{jk}^i X^k$$

satisfies the condition

$$\nabla_k (V_X)_j^i = \nabla_j (V_X)_k^i,$$

which is well-known in the Hamiltonian theory of systems of hydrodynamic type [8].

5.2 Integrability condition

In the flat case, we have seen that system (23) admits a set of solutions, given by the vector fields of the principal hierarchy. However, if ∇ is non-flat, existence of solutions for system (23) is not guaranteed; additional constraints have to be imposed on the curvature R of the connection ∇ .

Proposition 16 If X is a solution of (22), then the identity

$$Z \circ R(W, Y)(X) + W \circ R(Y, Z)(X) + Y \circ R(Z, W)(X) = 0, \quad (27)$$

holds for any choice of the vector fields (Y, W, Z) .

Proof. Condition (22) implies

$$\nabla_W (Z \circ \nabla_Y X - Y \circ \nabla_Z X) + \nabla_Y (W \circ \nabla_Z X - Z \circ \nabla_W X) + \nabla_Z (Y \circ \nabla_W X - W \circ \nabla_Y X) = 0.$$

Using the symmetry condition (17) written in the form

$$\nabla_Y (X \circ Z) - \nabla_X (Y \circ Z) + Y \circ \nabla_X Z - X \circ \nabla_Y Z - [Y, X] \circ Z = 0$$

we obtain identity (27). \square

Condition (27) must be satisfied for *any* solution X of the system (23). Since we are looking for a family of vector fields satisfying (23), it is natural to require that (27) holds true for an *arbitrary* vector field X .

Definition 17 An F -manifold with compatible connection is a manifold endowed with an associative commutative multiplicative structure given by a $(1, 2)$ -tensor field c and a torsionless connection ∇ satisfying condition (18) and condition

$$Z \circ R(W, Y)(X) + W \circ R(Y, Z)(X) + Y \circ R(Z, W)(X) = 0, \quad (28)$$

for any choice of the vector fields (X, Y, W, Z) . In local coordinates this means that

$$R_{lmi}^k c_{pk}^n + R_{lip}^k c_{mk}^n + R_{lpm}^k c_{ik}^n = 0. \quad (29)$$

Remark 18 An equivalent form of condition (28) can be easily obtained using the (second) Bianchi identity for the deformed connection

$$\tilde{\nabla}_X Y = \nabla_X Y + zX \circ Y, \quad z \in \mathbb{C},$$

where X and Y are arbitrary vector fields. Indeed, due to associativity and symmetry condition (17), the Riemann tensor of this connection does not depend on z [24]. Using this fact it is easy to see that the Bianchi identity reduces to

$$\begin{aligned} 0 &= \tilde{\nabla}_X R(Y, Z)(W) + \tilde{\nabla}_Z R(X, Y)(W) + \tilde{\nabla}_Y R(Z, X)(W) \\ &= X \circ R(Y, Z)(W) + Z \circ R(X, Y)(W) + Y \circ R(Z, X)(W) \\ &\quad - R(Y, Z)(X \circ W) - R(X, Y)(Z \circ W) - R(Z, X)(Y \circ W) \end{aligned}$$

for any choice of the vector fields (X, Y, W, Z) . Hence, condition (28) is equivalent to

$$R(Y, Z)(X \circ W) + R(X, Y)(Z \circ W) + R(Z, X)(Y \circ W) = 0,$$

for every (X, Y, W, Z) .

From now on we will assume the existence of canonical coordinates (r^1, \dots, r^n) , discussing the meaning of condition (29) under this additional assumption.

Proposition 19 In canonical coordinates, system (23) reduces to

$$\partial_k v^i = \Gamma_{ki}^i (v^k - v^i), \quad i \neq k, \quad (30)$$

where v^i are the components of X in such coordinates.

Proof. Writing (23) in canonical coordinates, we get

$$\delta_j^i (\partial_k v^i + \Gamma_{kl}^i v^l) = \delta_k^i (\partial_j v^i + \Gamma_{jl}^i v^l).$$

In the case $i = j \neq k$, using the identities

$$\Gamma_{kk}^i = -\Gamma_{ki}^i \quad (31)$$

and

$$\Gamma_{kl}^i = 0, \quad i \neq k \neq l \neq i, \quad (32)$$

which follow from (17), we obtain system (30). The remaining conditions give no further constraints. \square

Remark 20 We recall that, in canonical coordinates, the components of the vector field X coincide with the characteristic velocities of the associated system of hydrodynamic type:

$$r_t^i = c_{jk}^i v^k r_x^j = v^i r_x^i, \quad i = 1, \dots, n.$$

Compatibility conditions of system (30) are well-known in the literature [25], and are given by the following conditions:

$$\partial_i \Gamma_{mk}^k - \partial_m \Gamma_{ik}^k = 0, \quad (33)$$

$$\partial_i \Gamma_{km}^k - \Gamma_{km}^k \Gamma_{im}^m + \Gamma_{ik}^k \Gamma_{km}^k - \Gamma_{ik}^k \Gamma_{im}^i = 0, \quad (34)$$

for pairwise distinct indices k, i, m .

Proposition 21 Condition (29) is equivalent to conditions (33) and (34).

Proof. In canonical coordinates, condition (29) reads

$$\begin{aligned} R_{lmi}^k c_{pk}^n + R_{lip}^k c_{mk}^n + R_{lpm}^k c_{ik}^n = \\ R_{lmi}^k \delta_p^n \delta_k^n + R_{lip}^k \delta_m^n \delta_k^n + R_{lpm}^k \delta_i^n \delta_k^n = \\ R_{lmi}^n \delta_p^n + R_{lip}^n \delta_m^n + R_{lpm}^n \delta_i^n = 0. \end{aligned}$$

If all the indices m, i, p, n are distinct the above condition is trivially satisfied. Let us consider the case $n = p$ (the case $n \neq p$ can be treated in the same way and does not add further condition). If $n = i$, we obtain

$$R_{lmn}^n + R_{lnm}^n + \delta_m^n R_{lnn}^n = 0,$$

that is satisfied due to the skew-symmetry of the Riemann tensor with respect to the second and third lower indices. The same if $n = m$. For $n \neq i, m$, we obtain

$$R_{nmi}^n = 0, \quad (35)$$

if $l = n$ and

$$R_{lmi}^n = 0, \quad (36)$$

if $l \neq n$. Since, due to (31), the components of the Riemann tensor vanish if all the indices are distinct, condition (36) reduces to

$$R_{mni}^n = 0, \quad n \neq m \neq i \neq n. \quad (37)$$

Finally, using (31) and (32), it is easy to check that conditions (35) and (37) are equivalent to conditions (33) and (34) respectively. This proves the proposition. \square

Remark 22 *If the compatibility conditions (33) and (34) are satisfied, the general solution of the system (30) depends on n arbitrary functions of a single variable. Moreover, due to (33), any solution (v^1, \dots, v^n) of (30) satisfies the condition*

$$\partial_k \left(\frac{\partial_j v^i}{v^j - v^i} \right) - \partial_j \left(\frac{\partial_k v^i}{v^k - v^i} \right) = 0, \quad i \neq j \neq k \neq i, \quad (38)$$

known in literature as semi-Hamiltonian property [25]. An invariant and highly non trivial formulation of such a property has been found in [22].

Due to the above remark, under the assumption of existence of canonical coordinates we have a set of solutions of (30) leading to a family of commuting systems of hydrodynamic type, depending on n arbitrary functions. This result shows the deep relation between F -manifold with compatible connection (Definition 17) and integrable systems of PDEs.

6 Riemannian F -manifolds and Egorov metrics

In this section we consider the special case where the connection ∇ is a metric connection. This assumption plays an important role in the Hamiltonian theory of systems of hydrodynamic type (see for instance [3, 21, 23] and references therein), as well as in the theory of Frobenius manifolds [4, 5].

Definition 23 *A Riemannian F -manifold is an F -manifold with a compatible connection ∇ satisfying the following additional conditions:*

1. *The connection is metric:*

$$\nabla g = 0 .$$

2. *The inner product $\langle \cdot, \cdot \rangle$ defined by the metric g is invariant with respect to the product \circ :*

$$\langle X \circ Y, Z \rangle = \langle X, Y \circ Z \rangle . \quad (39)$$

In local coordinates, condition (39) reads

$$g_{iq} c_{lp}^q = g_{lq} c_{ip}^q, \quad \text{or} \quad g^{iq} c_{qp}^l = g^{lq} c_{qp}^i, \quad (40)$$

where g_{ij} and g^{ij} are respectively the covariant and the contravariant components of the metric g .

If there exist canonical coordinates, the metric g entering the definition of Riemannian F -manifold is an Egorov metric. Let us recall the definition of this special class of metrics.

Definition 24 *A metric is called Egorov if there exist coordinates (r^1, \dots, r^n) such that it is diagonal and potential:*

$$g_{ij} = \delta_j^i g_{ii}(r^1, \dots, r^n) = \delta_j^i \partial_i F,$$

for a certain function F .

Now, if we assume the existence of canonical coordinates, condition (40) tells us that the metric g is diagonal in such coordinates, while condition (32)—which follows from (17)—implies that the metric is potential. Therefore, g is an Egorov metric. Conversely, given an Egorov metric g whose curvature tensor satisfies condition (37), we can locally construct a Riemannian F -manifold. More precisely, let (r^1, \dots, r^n) be the coordinates where g is diagonal and potential. Then, the metric g and the structure constants

$$c_{jk}^i(r) = \delta_j^i \delta_k^i$$

endow the open set where the coordinates (r^1, \dots, r^n) are defined with the structure of a Riemannian F -manifold.

We point out that condition (29) is far from being trivial. Indeed, using the above remark, it is easy to construct examples of metrics satisfying properties (39) and (17). Much more difficult is the problem of finding Egorov metrics which satisfy also condition (29), since the potential has to fulfill (37). However, there exists an important class of metrics, appearing in the Hamiltonian theory of integrable hierarchies of hydrodynamic type (not necessarily of Egorov type) whose curvature satisfies (29). These are the metrics whose Riemann tensor admits “a quadratic expansion” in terms of the flows of the hierarchy [9, 10]:

$$u_{t_\alpha}^i = c_{jk}^i X_{(\alpha)}^k u_x^j, \quad i = 1, \dots, n.$$

This means that

$$R_{mi}^{sk} = (c_{ml}^s c_{iq}^k - c_{il}^s c_{mq}^k) \sum_{\alpha} \epsilon_{\alpha} X_{(\alpha)}^l X_{(\alpha)}^q, \quad \epsilon_{\alpha} = \pm 1, \quad (41)$$

where the index α can take value on a finite or infinite—even continuous—set.

Proposition 25 *Suppose that ∇ is the Levi-Civita connection of a metric g , and that its curvature satisfies condition (41). In this case, condition (29) is automatically satisfied.*

Proof. We have that

$$\begin{aligned} & R_{mi}^{sk} c_{pk}^n + R_{ip}^{sk} c_{mk}^n + R_{pm}^{sk} c_{ik}^n = \\ & \sum_{\alpha} \epsilon_{\alpha} [(c_{mr}^s c_{iq}^k - c_{ir}^s c_{mq}^k) c_{pk}^n + (c_{ir}^s c_{pq}^k - c_{pr}^s c_{iq}^k) c_{mk}^n + (c_{pr}^s c_{mq}^k - c_{mr}^s c_{pq}^k) c_{ik}^n] X_{(\alpha)}^r X_{(\alpha)}^q = \\ & \sum_{\alpha} \epsilon_{\alpha} [(c_{iq}^k c_{pk}^n - c_{pq}^k c_{ik}^n) c_{mr}^s + (c_{pq}^k c_{mk}^n - c_{mq}^k c_{pk}^n) c_{ir}^s + (c_{mq}^k c_{ik}^n - c_{iq}^k c_{mk}^n) c_{pr}^s] X_{(\alpha)}^r X_{(\alpha)}^q, \end{aligned}$$

which vanishes due to associativity. \square

Remark 26 *If the functions*

$$g^{lq} := \sum_{\alpha} \epsilon_{\alpha} X_{(\alpha)}^l X_{(\alpha)}^q$$

define the contravariant components of a metric satisfying condition (40), then the operator

$$\sum_{\alpha} \epsilon_{\alpha} (w_{\alpha})_k^i u_x^k \left(\frac{d}{dx} \right)^{-1} (w_{\alpha})_h^j u_x^h, \quad (w_{\alpha})_j^i := c_{jk}^i X_{(\alpha)}^k$$

is a purely nonlocal Poisson operator (see [13] for details).

7 An example: reductions of the dispersionless KP hierarchy

In this section we will consider a class of Riemannian F -manifolds associated with a well-known class of hydrodynamic type systems: the reductions of the dispersionless KP hierarchy. For a generic reduction, the metric will be non-flat.

The dispersionless KP (or dKP) hierarchy can be defined by introducing the formal series

$$\lambda = p + \sum_{k=0}^{\infty} \frac{A^k}{p^{k+1}}, \quad (42)$$

which has to satisfy the following dispersionless Lax equations

$$\lambda_{t_n} = \left\{ \lambda, \frac{1}{n} (\lambda^n)_+ \right\}.$$

Here $\{f, g\} = \partial_x f \partial_p g - \partial_p f \partial_x g$ denotes the canonical Poisson bracket, and $(\cdot)_+$ is the polynomial part of the argument. For simplicity, we will consider here only the second flow ($n = 2$); all other flows of the hierarchy can be treated in the same way. For the second flow, we have

$$\lambda_{t_2} = \left\{ \lambda, \frac{1}{2} p^2 + A^0 \right\} = p \lambda_x - A_x^0 \lambda_p, \quad (43)$$

or, explicitly in terms of the variables A^k ,

$$A_{t_2}^k = A_x^{k+1} + k A^{k-1} A_x^0, \quad k = 0, 1, 2, \dots \quad (44)$$

This last system is also known in the literature as Benney chain [1]; its Lax representation (43) appeared for the first time in [19]. An n -component reduction of the Benney chain is a restriction of the infinite dimensional system (44) to a suitable n -dimensional submanifold, that is

$$A^k = A^k(u^1, \dots, u^n), \quad k = 0, 1, \dots$$

The reduced systems are systems of hydrodynamic type in the variables (u^1, \dots, u^n) that parametrize the submanifold:

$$u_t^i = v_j^i(u) u_x^j, \quad i = 1, \dots, n.$$

Reductions of the Benney system were introduced in [14], and there it was proved that such systems are diagonalizable and integrable via the generalized hodograph transformation [25]. Clearly, in the case of a reduction, the coefficients of the series (42) depend on the Riemann invariants (r^1, \dots, r^n) and the series can be thought as the asymptotic expansion for $p \rightarrow \infty$ of a suitable function $\lambda(p, r^1, \dots, r^n)$ depending piecewise analytically on the parameter p . It turns out [14, 15] that such a function satisfies a system of chordal Loewner equations,

$$\frac{\partial \lambda}{\partial r^i} = \frac{\partial_i A^0}{p - v^i} \lambda_p, \quad i = 1, \dots, n, \quad (45)$$

describing families of conformal maps (with respect to p) in the complex upper half plane. The analytic properties of λ characterize the reduction. More precisely, in the case of an n -reduction the associated function λ possesses n distinct critical points on the real axis; these are the characteristic velocities v^i of the reduced system, that is,

$$\lambda_p(v^i) := \frac{\partial \lambda}{\partial p}(v^i) = 0, \quad i = 1, \dots, n,$$

and the corresponding critical values can be chosen as Riemann invariants. Compatibility conditions of the Loewner system (45) are of the form

$$\begin{aligned} \partial_i v^j &= \frac{\partial_i A^0}{v^i - v^j} \\ & \quad i \neq j, \\ \partial_{ij}^2 A^0 &= \frac{2\partial_i A^0 \partial_j A^0}{(v^i - v^j)^2} \end{aligned}$$

and were found by Gibbons and Tsarev [15]. Thus, every reduction of the Benney chain is described by a particular solution of the Loewner system (45).

Starting from the function λ , we will show now how to give to the manifold parametrized by the Riemann invariants (r^1, \dots, r^n) , a structure of F -manifold with a compatible connection—in general non-flat. In order to do this, we define a metric

$$g(\partial, \partial') = \sum_{i=1}^n \operatorname{res}_{p=v^i} \left(\frac{\partial \lambda(p) \partial' \lambda(p)}{\lambda_p} dp \right), \quad (46)$$

and structure constants

$$c(\partial, \partial', \partial'') = \sum_{i=1}^n \operatorname{res}_{p=v^i} \left(\frac{\partial \lambda(p) \partial' \lambda(p) \partial'' \lambda(p)}{\lambda_p} dp \right), \quad (47)$$

where $\partial, \partial', \partial''$ are arbitrary tangent vectors on the manifold. In the coordinates (r^1, \dots, r^n) , and making use of the Loewner equations (45), the metric takes the diagonal form

$$\begin{aligned} g \left(\frac{\partial}{\partial r^i}, \frac{\partial}{\partial r^j} \right) &= \sum_{i=1}^n \operatorname{res}_{p=v^i} \left(\frac{\partial \lambda}{\partial r^i} \frac{\partial \lambda}{\partial r^j} \frac{dp}{\lambda_p} \right) = \sum_{i=1}^n \operatorname{res}_{p=v^i} \left(\frac{\partial_i A^0 \partial_j A^0 \lambda_p dp}{(p - v^i)(p - v^j)} \right) \\ &= \partial_i A^0 \partial_j A^0 \lambda_{pp}(v^i) \delta_{ij} = \partial_i A^0 \delta_{ij}, \end{aligned}$$

where we used the fact [12] that

$$\lambda_{pp}(v^i) = \frac{1}{\partial_i A^0}.$$

In particular, the metric is Egorov. Moreover, a similar calculation for the structure constants gives

$$\begin{aligned} c\left(\frac{\partial}{\partial r^i}, \frac{\partial}{\partial r^j}, \frac{\partial}{\partial r^k}\right) &= \sum_{i=1}^n \operatorname{res}_{p=v^i} \left(\frac{\partial \lambda}{\partial r^i} \frac{\partial \lambda}{\partial r^j} \frac{\partial \lambda}{\partial r^k} \frac{dp}{\lambda_p} \right) = \sum_{i=1}^n \operatorname{res}_{p=v^i} \left(\frac{\partial_i A^0 \partial_j A^0 \partial_k A^0 (\lambda_p)^2 dp}{(p-v^i)(p-v^j)(p-v^k)} \right) \\ &= \partial_i A^0 \partial_j A^0 \partial_k A^0 (\lambda_{pp}(v^i))^2 \delta_{ij} \delta_{ik} = \partial_i A^0 \delta_{ij} \delta_{ik}, \end{aligned}$$

and from this it follows that

$$\frac{\partial}{\partial r^i} \circ \frac{\partial}{\partial r^j} = \delta_{ij} \frac{\partial}{\partial r^i},$$

namely (r^1, \dots, r^n) are canonical coordinates for the algebra.

Remark 27 *The metric (46) and the structure constants (47) were introduced for the first time by Dubrovin in [4], in the particular case of the Gelfand-Dikii reductions of the dKP hierarchy, where the function λ is a polynomial in p . The same metric and constants were also used by Chang [2] and Ferguson and Strachan [11], for the study of reductions where λ is rational or logarithmic. We remark that in all these examples the metric considered turns out to be flat.*

We have now to prove that the metric and the structure constants defined in this way are compatible, namely that conditions (17) and (29) are satisfied. As regard condition (17)—due to the results of Section 6—it is sufficient to note that the metric (46) is Egorov. On the other hand, for condition (29), we only have to recall the result of [12], where the curvature tensor of the metric (46) has been shown to possess the following quadratic expansion:

$$R_{ij}^{ij} = \frac{1}{2\pi i} \int_C w^i(\lambda) w^j(\lambda) d\lambda, \quad w^i(\lambda) = \frac{\frac{\partial p}{\partial \lambda}}{(p(\lambda) - v^i)^2},$$

where $p(\lambda) = \lambda^{-1}(p)$ is the inverse of λ with respect to p , and C is a suitable contour on the complex λ -plane. Due to Proposition 25, the existence of a quadratic expansion of the curvature implies that condition (29) is satisfied. Alternatively, such a condition follows from the well-known fact that the characteristic velocities v^i —which satisfy condition (30)—satisfy the semi-Hamiltonian condition (38).

Remark 28 *A similar construction can be done using instead of the metric (46), one of the metrics*

$$g\left(\frac{\partial}{\partial r^i}, \frac{\partial}{\partial r^j}\right) = \sum_{i=1}^n \operatorname{res}_{p=v^i} \varphi_i(r^i) \left(\frac{\partial \lambda}{\partial r^i} \frac{\partial \lambda}{\partial r^j} \frac{dp}{\lambda_p} \right) \quad (48)$$

where φ_i are arbitrary functions of a single variable, and defining the corresponding structure constants as

$$c\left(\frac{\partial}{\partial r^i}, \frac{\partial}{\partial r^j}, \frac{\partial}{\partial r^k}\right) = \sum_{i=1}^n \operatorname{res}_{p=v^i} (\varphi_i(r^i))^2 \left(\frac{\partial \lambda}{\partial r^i} \frac{\partial \lambda}{\partial r^j} \frac{\partial \lambda}{\partial r^k} \frac{dp}{\lambda_p} \right). \quad (49)$$

If all the functions φ_i are different from zero, it turns out that the structure constants (49) admit canonical coordinates. Moreover, in such coordinates, the metric (48) is potential. In this way, repeating the construction described in this section, one defines, for any choice of the functions φ_i , a new structure of F -manifold with compatible connection on the same manifold. Notice that in case of reductions related to Frobenius manifolds, such as the Zakharov and the Gel'fand-Dikii reductions [4, 7], one of the metrics (48) is the intersection form of the Frobenius manifold. Using this metric, the construction above reduces to the Dubrovin's duality of the theory of Frobenius manifolds [6].

Acknowledgments

We thank Maxim Pavlov for stimulating discussions. M.P. and A.R. would like to thank the Department *Matematica e Applicazioni* of the Milano-Bicocca University for the hospitality. Support of A.R. through the ESF grant MISGAM 2265 for his visit to the Milano-Bicocca department is gratefully acknowledged.

References

- [1] D.J. Benney, *Some properties of long nonlinear waves*, Stud. Appl. Math. **52** (1973), 45–50.
- [2] Jen-Hsu Chang, *On the waterbag model of the dispersionless KP hierarchy (II)*, J. Phys. A: Math. Theor. **40** (2007), 12973–12985.
- [3] B.A. Dubrovin, *On the differential geometry of strongly integrable systems of hydrodynamics type*, (Russian) Funktsional. Anal. i Prilozhen. **24** (1990), no. 4, 25–30, 96; translation in Funct. Anal. Appl. **24** (1990), no. 4, 280–285 (1991).
- [4] B.A. Dubrovin, *Geometry of 2D topological field theories*, in: Integrable Systems and Quantum Groups, Montecatini Terme, 1993. Editors: M. Francaviglia, S. Greco. Springer Lecture Notes in Math. **1620** (1996), pp. 120–348.
- [5] B.A. Dubrovin, *Flat pencils of metrics and Frobenius manifolds*, in: Integrable systems and algebraic geometry (Kobe/Kyoto,1997) World Sci. Publ., River Edge, NJ **1620** (1998), pp. 47–72.
- [6] B.A. Dubrovin, *On almost duality for Frobenius manifolds*, in: Geometry, topology, and mathematical physics, Amer. Math. Soc. Transl. Ser. 2, 212, Amer. Math. Soc., Providence, RI, 2004, pp. 75–132.
- [7] B.A. Dubrovin, S.Q. Liu, Y. Zhang, *Frobenius manifolds and central invariants for the Drinfeld-Sokolov biHamiltonian structures*, Adv. Math. **219** (2008), no. 3, 780–837.

- [8] B.A. Dubrovin, S.P. Novikov, *Hydrodynamics of weakly deformed soliton lattices. Differential geometry and Hamiltonian theory*, Uspekhi Mat. Nauk **44** (1989), 29–98. English translation in Russ. Math. Surveys **44** (1989), 35–124.
- [9] E.V. Ferapontov, *Differential geometry of nonlocal Hamiltonian operators of hydrodynamic type*, Funct. Anal. Appl. **25** (1991), no. 3, 195–204.
- [10] E.V. Ferapontov, O.I. Mokhov, *Nonlocal Hamiltonian operators of hydrodynamic type that are connected with metrics of constant curvature*, Russ. Math. Surv. **45** (1990), no. 3, 218–219.
- [11] J.T. Ferguson, I.A.B. Strachan, *Logarithmic deformations of the rational superpotential/Landau-Ginzburg construction of solutions of the WDVV equations*, Comm. Math. Phys. **280** (2008), 1–25.
- [12] J. Gibbons, P. Lorenzoni, A. Raimondo, *Hamiltonian structure of reductions of the Benney system*, Comm. Math. Phys. **287** (2009), 291–322,
- [13] J. Gibbons, P. Lorenzoni, A. Raimondo, *Purely nonlocal Hamiltonian formalism for systems of hydrodynamic type*, arXiv:0812.3317.
- [14] J. Gibbons, S.P. Tsarev, *Reductions of the Benney equations*, Phys. Lett. A **211** (1996), no. 1, 19–24.
- [15] J. Gibbons, S.P. Tsarev, *Conformal maps and reductions of the Benney equations*, Phys. Lett. A **258** (1999), no. 4–6, 263–271.
- [16] C. Hertling, *Multiplication on the tangent bundle*, arXiv:math/9910116
- [17] C. Hertling, Y. Manin, *Weak Frobenius manifolds*, Internat. Math. Res. Notices **1999**, no. 6, 277–286.
- [18] B.G. Konopelchenko, F. Magri, *Coisotropic deformations of associative algebras and dispersionless integrable hierarchies*, Comm. Math. Phys. **274** (2007), 627–658.
- [19] D. Lebedev, Y. Manin, *Conservation laws and Lax representation of Benney’s long wave equations*, Phys. Lett. A **74** (1979), 154–156.
- [20] Y. Manin, *F-manifolds with flat structure and Dubrovin’s duality*, Adv. Math. **198** (2005), no. 1, 5–26.
- [21] M.V. Pavlov, *Integrability of Egorov systems of hydrodynamic type*, (Russian) Teoret. Mat. Fiz. **150** (2007), no. 2, 263–285; translation in Theoret. and Math. Phys. **150** (2007), no. 2, 225–243.
- [22] M.V. Pavlov, S.I. Svinolupov, R.A. Sharipov, *An invariant criterion for hydrodynamic integrability*, (Russian) Funktsional. Anal. i Prilozhen. **30** (1996), no. 1, 18–29, 96; translation in Funct. Anal. Appl. **30** (1996), no. 1, 15–22.

- [23] M.V. Pavlov, S.P. Tsarev, *Tri-Hamiltonian structures of Egorov systems of hydrodynamic type*, (Russian) Funktsional. Anal. i Prilozhen. **37** (2003), no. 1, 38–54.
- [24] I.A.B. Strachan, *Frobenius manifolds: natural submanifolds and induced bi-Hamiltonian structures*, Differential Geom. Appl. **20** (2004), no. 1, 67–99.
- [25] S.P. Tsarev, *The geometry of Hamiltonian systems of hydrodynamic type. The generalised hodograph transform*, USSR Izv. **37** (1991) 397–419.
- [26] V.E. Zakharov, *Benney equations and quasiclassical approximation in the inverse problem*, Funktsional. Anal. i Prilozhen **14** (1980), 15–24.