

Asymptotic Behavior and Decay Estimates of the Solutions for a Nonlinear Parabolic Equation with Exponential Nonlinearity

Giulia FURIOLI

Università di Bergamo, Italy

Tatsuki KAWAKAMI

Osaka Prefecture University, Japan

Bernhard RUF

Università di Milano, Italy

Elide TERRANEO

Università di Milano, Italy

Abstract

We consider a nonlinear parabolic equation with an exponential nonlinearity which is critical with respect to the growth of the nonlinearity and the regularity of the initial data. After showing the equivalence of the notions of weak and mild solutions, we derive decay estimates and the asymptotic behavior for small global-in-time solutions.

Keywords: asymptotic behavior, nonlinear heat equation, fractional diffusion equation, exponential nonlinearity, Orlicz space

1 Introduction

We consider the Cauchy problem for the semilinear heat equation

$$\begin{cases} \partial_t u = \Delta u + f(u), & t > 0, x \in \mathbb{R}^n, \\ u(0, x) = \varphi(x), & x \in \mathbb{R}^n, \end{cases} \quad (1.1)$$

where $n \geq 1$, $u(t, x) : \mathbb{R}^+ \times \mathbb{R}^n \rightarrow \mathbb{R}$ is the unknown function, $\partial_t = \partial/\partial t$, $f(u)$ contains the nonlinearity and φ is the initial data. This paper is concerned with the asymptotic behavior and decay estimates of the solutions of (1.1) in certain limiting cases which are critical with respect to the growth of the nonlinearity and the regularity of the initial data.

Before introducing the subject of this paper, let us recall some related known results.

The polynomial case. The case of power nonlinearities $f(u) = |u|^{p-1}u$ with $p > 1$, that is

$$\begin{cases} \partial_t u = \Delta u + |u|^{p-1}u, & t > 0, x \in \mathbb{R}^n, \\ u(0, x) = \varphi(x), & x \in \mathbb{R}^n, \end{cases} \quad (1.2)$$

has been extensively studied since the pioneering works by Fujita [6] and Weissler [22, 23]. It is well-known that the problem (1.2) satisfies a scale invariance property. In fact, for $\lambda \in \mathbb{R}_+$, if u is a solution of (1.2), then

$$u_\lambda(t, x) := \lambda^{\frac{2}{p-1}} u(\lambda^2 t, \lambda x) \quad (1.3)$$

is also a solution of (1.2) with initial data $\varphi_\lambda(x) := \lambda^{2/(p-1)} \varphi(\lambda x)$. So, all function spaces invariant with respect to the scaling transformation (1.3) play a fundamental role in the study of the Cauchy problem (1.2). In the framework of Lebesgue spaces, we can easily show that the norm of the space $L^q(\mathbb{R}^n)$ is invariant with respect to (1.3) if and only if $q = q_c$ with $q_c = n(p-1)/2$, and it is well-known that, given the nonlinearity $|u|^{p-1}u$, the Lebesgue space $L^{q_c}(\mathbb{R}^n)$ plays the role of *critical space* for the well-posedness of (1.2) (see e.g. [2, 7, 22, 23]).

Indeed, for any $q \geq q_c$ and $q > 1$, or $q > q_c$ and $q \geq 1$ (*subcritical case*), and for any initial data $\varphi \in L^q(\mathbb{R}^n)$ there exists a local (in time) solution of the Cauchy problem (1.2). On the other hand, for initial data belonging to $L^q(\mathbb{R}^n)$ with $1 < q < q_c$ (*supercritical case*), Weissler [22] and Brezis-Cazenave [2] indicate that for certain $\varphi \in L^q(\mathbb{R}^n)$ there exists no local (in time) solution in any reasonable sense.

We can also state the previous results from a different point of view. Given any initial data in the Lebesgue space L^q , $1 < q < +\infty$, the Cauchy problem (1.2) is well-posed if and only if the power nonlinearity p is smaller than or equal to the critical value $p_c = 1 + (2q)/n$. Moreover, the *critical case*, which is defined equivalently by $q = q_c$ or $p = p_c$, is the only case for which global (in time) existence can be established for small initial data.

The same polynomial nonlinearity has also been considered for the Schrödinger equation

$$\begin{cases} i\partial_t u + \Delta u = |u|^{p-1}u, & t > 0, x \in \mathbb{R}^n, \\ u(0, x) = \varphi(x), & x \in \mathbb{R}^n. \end{cases} \quad (1.4)$$

Here $u : \mathbb{R}^+ \times \mathbb{R}^n \rightarrow \mathbb{C}$ and $\varphi \in H^s(\mathbb{R}^n)$ with $0 \leq s < n/2$. Also for problem (1.4) the scaling invariance (1.3) holds, and the scaling argument indicates the critical Sobolev space and exponent for the nonlinearity $|u|^{p-1}u$: for any fixed $p > 1$, the Sobolev space $H^{s_c}(\mathbb{R}^n)$ with $s_c = n/2 - 2/(p-1)$ plays the role of *critical space* for the well-posedness of (1.4) (see e.g. [3]). Indeed, for initial data $\varphi \in H^s(\mathbb{R}^n)$ with $s \geq s_c$ (*subcritical case*) Cazenave and Weissler [3] proved that there exists a local (in time) solution in $H^s(\mathbb{R}^n)$.

Equivalently, for any initial data in the Sobolev space $H^s(\mathbb{R}^n)$, $0 \leq s < n/2$, local existence for the Cauchy problem (1.4) holds true under the condition $p \leq \tilde{p}_c = 1 + 4/(n-2s)$, namely for any power nonlinearity smaller than or equal to the critical nonlinearity \tilde{p}_c .

Moreover, the *critical case*, which is defined equivalently by $s = s_c$ or $p = \tilde{p}_c$, is the only one for which global (in time) existence can be established for small initial data.

The case $s_c = n/2$ is a *limiting case*, since for $s > n/2$ the Sobolev space $H^s(\mathbb{R}^n)$ embeds into $L^\infty(\mathbb{R}^n)$, and no specific behavior for the nonlinearity is required for the existence of local-in-time solutions of (1.4) in the H^s framework.

The limiting case. Let us now consider the *limiting critical case* $H^{n/2}(\mathbb{R}^n)$; then

(i) any power nonlinearity is subcritical, since $H^{n/2}(\mathbb{R}^n)$ embeds into any $L^p(\mathbb{R}^n)$ space,

(ii) $H^{n/2}(\mathbb{R}^n)$ does not embed into $L^\infty(\mathbb{R}^n)$, and by Trudinger's inequality [20] one knows that $H^{n/2}(\mathbb{R}^n)$ embeds into the Orlicz space $L_\varphi(\mathbb{R}^n)$, with growth function (N -function) $\varphi(s) = e^{s^2} - 1$.

For this limiting case, Nakamura and Ozawa [15] considered the Cauchy problem

$$\begin{cases} i\partial_t u + \Delta u = f(u), & t > 0, x \in \mathbb{R}^n, \\ u(0) = \varphi \in H^{\frac{n}{2}}(\mathbb{R}^n), \end{cases} \quad (1.5)$$

with exponential nonlinearity of asymptotic growth $f(u) \sim e^{u^2}$ and with a vanishing behavior at the origin. They proved the existence of global-in-time solutions of (1.5) for initial data with small norm in $H^{n/2}(\mathbb{R}^n)$. In view of the Trudinger inequality the growth rate of $f(u)$ at infinity seems to be optimal for initial data in $H^{n/2}(\mathbb{R}^n)$.

It is well-known that there is a correspondence between the $H^s(\mathbb{R}^n)$ theory for the nonlinear Schrödinger equation (1.4) and the $L^q(\mathbb{R}^n)$ theory for the semilinear heat equation (1.2) (see e.g. [3]). Indeed, since the Sobolev space $H^s(\mathbb{R}^n)$ embeds into $L^q(\mathbb{R}^n)$ with $q = (2n)/(n - 2s)$, for $0 \leq s < n/2$, it holds that $p_c = 1 + (2q)/n = 1 + 4/(n - 2s) = \tilde{p}_c$. This means that the critical nonlinearity $|u|^{p_c-1}u$ associated to (1.2) in the $L^q(\mathbb{R}^n)$ framework corresponds to the critical one associated to (1.4) in the $H^s(\mathbb{R}^n)$ framework.

As a natural analogy to the results of [15], the third and fourth authors of this paper [19] and Ioku [8, 9] considered the Cauchy problem (1.1) with a nonlinearity of the form

$$f(u) = |u|^{\frac{4}{n}} u e^{u^2} \quad (1.6)$$

and initial data φ belonging to the Orlicz space $\exp L^2$ defined as

$$\exp L^2(\mathbb{R}^n) := \left\{ u \in L^1_{loc}(\mathbb{R}^n), \exists \lambda > 0 : \int_{\mathbb{R}^n} \left(\exp \left(\frac{|u(x)|}{\lambda} \right)^2 - 1 \right) dx < \infty \right\}$$

(see also Definition 3.1). Clearly, no scaling invariance holds true for equation (1.1) with such a nonlinear term. Also, the growth e^{u^2} at infinity of the nonlinearity $f(u)$ seems to be optimal in the framework of the Orlicz space $\exp L^2$. In fact, if $f(u) \sim e^{|u|^r}$ with $r > 2$, then there exist positive initial data $\varphi \in \exp L^2$ (even with very small norm) such that there exists no nonnegative classical local-in-time solution of (1.1) (see [10]).

For the Cauchy problem (1.1) with (1.6), the authors of the papers [8, 9, 19] considered the corresponding integral equation

$$u(t) = e^{t\Delta} \varphi + \int_0^t e^{(t-s)\Delta} f(u(s)) ds. \quad (1.7)$$

As we will recall in Definition 2.2, u is called a *mild* solution of (1.1) if u is a solution of the integral equation (1.7) and $u(t) \xrightarrow[t \rightarrow 0]{} \varphi$ in the weak* topology (see (2.1)). Under this notion of solution one has

Proposition 1.1 [8, 9, 19] *Let $n \geq 1$ and $\varphi \in \exp L^2$. Suppose that f satisfies (1.6). Then there exists $\varepsilon = \varepsilon(n) > 0$ such that, if $\|\varphi\|_{\exp L^2} < \varepsilon$, then there exists a mild solution u of (1.1) satisfying*

$$u \in L^\infty(0, \infty; \exp L^2)$$

and

$$\sup_{t>0} \|u(t)\|_{\exp L^2} \leq 2\|\varphi\|_{\exp L^2}. \quad (1.8)$$

By these results one obtains the small data global existence of *mild* solutions of (1.1). However, as far as we know, there are no results which treat the correspondence between mild solutions and other notions of weak solutions, and which prove decay estimates and the asymptotic behavior of the solutions of (1.1).

In this paper, under a smallness assumption for the solution, we prove the equivalence between *mild* solutions and *weak* solutions of (1.1). Furthermore, under condition (1.8), we obtain decay estimates for the solutions in the following two cases

- $\varphi \in \exp L^2$ only (*singular case*), and
- $\varphi \in \exp L^2 \cap L^p(\mathbb{R}^n)$ with $p \in [1, 2)$ (*regular case*).

In particular, for the regular case $p = 1$, we show that global-in-time solutions with some suitable decay estimates behave asymptotically like suitable multiples of the heat kernel.

The paper is organized as follows. In Section 2 we state the main results. In Section 3 we give some preliminaries. Sections 4–7 are devoted to the proof of the theorems, respectively. In Section 8 we extend our theorems to fractional diffusion equations with general initial data.

Before closing this section we give some notations used in this paper. For any $p \in [1, \infty]$, let $\|\cdot\|_{L^p}$ be the usual norm of $L^p := L^p(\mathbb{R}^n)$; we write $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$; $\mathcal{D}(\Omega)$ denotes the space of C^∞ -functions with compact support in Ω , and $\mathcal{D}'(\Omega)$ the topological dual of $\mathcal{D}(\Omega)$. We denote by $\mathcal{S}'(\Omega)$ the space of tempered distributions. Throughout the present paper, C will denote a generic positive constant which may have different values also within the same line.

2 Main results

In this section we state the main results of this paper. Before presenting the main theorems we introduce the definition of *weak* and *mild* solutions.

Definition 2.1 (Weak solution) *Let $\varphi \in \exp L^2$, $T \in (0, \infty]$ and $u \in L^\infty(0, T; \exp L^2)$. We call u a weak solution of the Cauchy problem (1.1) in $(0, T) \times \mathbb{R}^n$ if u satisfies*

$$\partial_t u = \Delta u + f(u)$$

in the sense of distributions $\mathcal{D}'((0, T) \times \mathbb{R}^n)$ and $u(t) \xrightarrow[t \rightarrow 0]{} \varphi$ in weak topology.*

Definition 2.2 (Mild solution) *Let $\varphi \in \exp L^2$, $T \in (0, \infty]$ and $u \in L^\infty(0, T; \exp L^2)$. We call u a mild solution of the Cauchy problem (1.1) in $(0, T) \times \mathbb{R}^n$ if $u \in C((0, T); \exp L^2)$, u is a solution of the integral equation*

$$u(t) = e^{t\Delta}\varphi + \int_0^t e^{(t-s)\Delta} f(u(s)) ds$$

and $u(t) \xrightarrow[t \rightarrow 0]{} \varphi$ in the weak topology.*

We recall that $u(t) \xrightarrow[t \rightarrow 0]{} \varphi$ in weak* topology if and only if

$$\lim_{t \rightarrow 0} \int_{\mathbb{R}^n} (u(t, x)\psi(x) - \varphi(x)\psi(x)) dx = 0 \quad (2.1)$$

for every ψ belonging to the predual space of $\exp L^2$ (see Section 3).

Now we are ready to state the main results of this paper. First we show the equivalence between small *mild* solutions and small *weak* solutions of (1.1).

Proposition 2.1 *Let $\varphi \in \exp L^2$, $T \in (0, \infty]$ and $u \in L^\infty(0, T; \exp L^2)$. Then there exists a positive constant $\varepsilon > 0$ such that, if $\sup_{0 < t < T} \|u(t)\|_{\exp L^2} \leq \varepsilon$, then the following statements are equivalent:*

- (i) u is a weak solution of (1.1) in $(0, T) \times \mathbb{R}^n$;
- (ii) u is a mild solution of (1.1) in $(0, T) \times \mathbb{R}^n$.

Here and in the rest of the paper, for simplicity, we call u a solution of (1.1) if u is a mild solution of (1.1) in $(0, \infty) \times \mathbb{R}^n$.

Next we prove uniqueness of small solutions in $L^\infty((0, T); \exp L^2)$ for all dimensions $n \in \mathbb{N}$.

Proposition 2.2 *Let $n \geq 1$. There is $\varepsilon > 0$ such that if $\varphi \in \exp L^2(\mathbb{R}^n)$ with $\|\varphi\|_{\exp L^2} \leq \varepsilon$ and u, v are two solutions of (1.1) satisfying*

$$\sup_{t < T} \|u(t)\|_{\exp L^2} \leq 2\|\varphi\|_{\exp L^2}, \quad \sup_{t < T'} \|u(t)\|_{\exp L^2} \leq 2\|\varphi\|_{\exp L^2}, \quad (2.2)$$

then $u(t) = v(t)$ on $[0, \min(T, T')]$.

The following result provides a decay estimate of the small solution of (1.1) for the singular case, that is, $\varphi \in \exp L^2$ only.

Theorem 2.1 *Let $n \geq 1$ and $\varphi \in \exp L^2$ with $\varphi \geq 0$. Assume that there exists a unique positive solution u of (1.1) satisfying (1.8). Then there exist constants $\varepsilon = \varepsilon(n) > 0$ and $C = C(n) > 0$ such that, if $\|\varphi\|_{\exp L^2} < \varepsilon$, then the solution u satisfies*

$$\|u(t)\|_{L^q} \leq Ct^{-\frac{n}{2}(\frac{1}{2} - \frac{1}{q})} \|\varphi\|_{\exp L^2}, \quad t > 0, \quad (2.3)$$

for all $q \in [2, \infty]$.

Remark 2.1 (i) *By Propositions 1.1 and 2.2, if $\|\varphi\|_{\exp L^2}$ is small enough, then we can show that the assumption for the solution u is not empty.*

(ii) *We obtained the same decay estimate as the solution of the linear heat equation with initial data in L^2 .*

(iii) *For the lower dimensional case $1 \leq n \leq 4$, in the proof of Proposition 1.1, Ioku [8, 9] already obtained the decay estimate (2.3) for $1 + 4/n < q < 2 + 8/n$.*

(iv) *Due to the embedding $\exp L^2 \subset L^q$ for $2 \leq q < \infty$ (see Lemma 3.2), if $\varphi \in \exp L^2$ with $\|\varphi\|_{\exp L^2} < \varepsilon$ as in Proposition 1.1, then $u \in L^\infty(0, \infty; L^q)$ for all $2 \leq q < \infty$.*

Next we consider the regular case, namely $\varphi \in \exp L^2 \cap L^p$, $p \in [1, 2)$. The nonlinearity (1.6) satisfies $f(x) \sim x^{1+4/n}$ for $x \rightarrow 0^+$. So, if $u \in L^\infty(0, \infty; L^q)$ for $q \geq 2$, then the nonlinear term $f(u)$ belongs to L^p for $p \geq (2n)/(n+4)$. In the lower dimensional case $1 \leq n \leq 4$, this means that $f(u) \in L^p$ for all $p \geq 1$ but for the higher dimensional case $n \geq 5$ this implies a true constraint. This is the reason why in the next theorems we have to introduce some parameters p_* , p_1 (and p_2 in Lemma 6.1) which are meaningful only for the higher dimensional case. More specifically, in the higher dimensional case, we can prove two kinds of results about the decay estimate of solutions of (1.1). In the first one (Theorem 2.2), we only assume a control of the $\exp L^2$ norm of the initial data, allowing the L^p -norm of the same data to be large (see Remark 6.3). This mild assumption on the initial data entails a decay estimate of $\|u(t)\|_{L^q}$ only for $q \geq p_* > p$. In the second result (Theorem 2.3), under a stronger assumption, that is, a smallness assumption for both the $\exp L^2$ and the L^p norm of the initial data, we can prove a better decay estimate on $\|u(t)\|_{L^q}$ for all $q \geq p$.

Theorem 2.2 *Assume the same conditions as in Theorem 2.1. Furthermore, assume $\varphi \in L^p$ for some $p \in [1, 2)$. Put*

$$p_* := \max \left\{ p, \frac{2n}{n+4} \right\}. \quad (2.4)$$

Then there exist positive constants $\varepsilon = \varepsilon(n)$, $C = C(n)$ and a positive function $F = F(n, p_, \|\varphi\|_{p_*})$ such that, if*

$$\|\varphi\|_{\exp L^2} < \min(\varepsilon, F(n, p_*, \|\varphi\|_{L^{p_*}})) \quad (2.5)$$

then the solution u satisfies

$$\|u(t)\|_{L^q} \leq C t^{-\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} \|\varphi\|_{\exp L^2 \cap L^{p_*}}, \quad t > 0, \quad (2.6)$$

for all $q \in [p_, \infty]$. In particular, if $p \in (p_1, 2)$, then*

$$\|u(t)\|_{L^q} = o \left(t^{-\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} \right), \quad t \rightarrow \infty, \quad (2.7)$$

where

$$p_1 := \max \left\{ 1, \frac{2n}{n+4} \right\}. \quad (2.8)$$

Theorem 2.3 *Assume the same conditions as in Theorem 2.2. Then there exists a positive constant $\delta = \delta(n)$ such that, if*

$$\max\{\|\varphi\|_{\exp L^2}, \|\varphi\|_{L^p}\} < \delta,$$

then (2.6) with $p_ = p$ holds for all $q \in [p, \infty]$. In particular, for all $q \in [p, \infty)$,*

$$\|u(t)\|_{L^q} \leq C(1+t)^{-\frac{n}{2} \left(\frac{1}{p} - \frac{1}{q} \right)} \|\varphi\|_{\exp L^2 \cap L^p}, \quad t > 0.$$

Furthermore, if $p \in (1, 2)$, then (2.7) with $p_ = p$ holds.*

Finally we address the question of the asymptotic behavior of solutions of (1.1) when $\varphi \in \exp L^2 \cap L^1$. We show that global-in-time solutions with suitable decay properties behave asymptotically like suitable multiples of the heat kernel.

Theorem 2.4 *Let $n \geq 1$, $\varphi \geq 0$ and $\varphi \in \exp L^2 \cap L^1$. Assume that $\|\varphi\|_{\exp L^2}$ is small enough. Furthermore, suppose that*

a) for $n \geq 1$,

$$\sup_{t>0} t^{\frac{n}{2}(1-\frac{1}{q})} \|u(t)\|_{L^q} < \infty, \quad q \in [1, \infty]; \quad (2.9)$$

b) for $n \geq 5$, assume moreover that there is $T^* > 0$ such that

$$\sup_{0 < t < T^*} \|u(t)\|_{L^{\frac{4}{n}+1}} < \infty. \quad (2.10)$$

Then there exists the limit

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}^n} u(t, x) dx = \int_{\mathbb{R}^n} \varphi(x) dx + \int_0^\infty \int_{\mathbb{R}^n} f(u(t, x)) dx dt := m_*$$

such that

$$\lim_{t \rightarrow \infty} t^{\frac{n}{2}(1-\frac{1}{q})} \|u(t) - m_* g(t)\|_{L^q} = 0, \quad q \in [1, \infty], \quad (2.11)$$

where $g(t, x) = G_2(t+1, x)$ is the heat kernel, that is,

$$G_2(t, x) = (4\pi t)^{-\frac{n}{2}} e^{-\frac{|x|^2}{4t}}.$$

Remark 2.2 (i) Conditions (2.9) and (2.10) are fulfilled for example under the hypotheses of Theorems 2.2 and 2.3.

(ii) The large time behavior and the decay properties of the solution of (1.2) have been widely studied (see, i.e., [4, 11, 12, 13, 14, 16, 21] and the references therein). In order to treat these topics for (1.2), the scale transformation (1.3) plays an important role. However, unfortunately, for the problem (1.1) with (1.6), we don't have such a scaling property. Instead of (1.3), applying the embedding $\exp L^2 \subset L^q$ for $2 \leq q < \infty$ and exploiting the uniqueness of the solution, we overcome this difficulty, and this is one of the novelties of this paper.

In the last section of the paper, we consider the Cauchy problem associated to the fractional diffusion equation

$$\begin{cases} \partial_t u + \mathcal{L}_\theta u = |u|^{\frac{r\theta}{n}} u e^{ur}, & t > 0, x \in \mathbb{R}^n, \\ u(0, x) = \varphi(x), & x \in \mathbb{R}^n, \end{cases} \quad (2.12)$$

where $0 < \theta \leq 2$, $r > 1$ and $\varphi \in \exp L^r$ (we will recall the definition of the Orlicz spaces $\exp L^r$ in Definition 3.1). Here the operator $\mathcal{L}_\theta := (-\Delta)^{\theta/2}$ is the fractional Laplacian defined by the Fourier transform \mathcal{F} as

$$\mathcal{L}_\theta \phi := \mathcal{F}^{-1} [|\xi|^\theta \mathcal{F}[\phi]]. \quad (2.13)$$

In the case $\theta = 2$, this generalization is suggested by the previous results for (1.2) in the framework of Sobolev spaces H_q^s . Indeed, for any $s \in \mathbb{R}$ and $1 < q < \infty$, the Cauchy problem (1.2) has also been studied for initial data $\varphi \in H_q^s$, where

$$H_q^s(\mathbb{R}^n) := \left\{ \psi \in \mathcal{S}'(\mathbb{R}^n) : (1 - \Delta)^{\frac{s}{2}} \psi \in L^q(\mathbb{R}^n) \right\}$$

(see, i.e., [18]). For the case $s < n/q$, a critical power nonlinearity appears in analogy with the theory in the Lebesgue framework. While for $s > n/q$, no growth condition is necessary to establish the existence of a solution, in the case $s = n/q$ any power nonlinearity is allowed and one wonders which is the optimal critical growth at infinity. By the Trudinger inequality in \mathbb{R}^n we have the embedding

$$H_q^s \hookrightarrow \exp L^\Phi.$$

Here $\exp L^\Phi$ is the Orlicz space defined by the convex function

$$\Phi(t) := \exp\left(t^{\frac{q}{q-1}}\right) - \sum_{j=0}^{k_0-1} \frac{t^{jq/(q-1)}}{j!},$$

where k_0 is the smallest integer satisfying $k_0 \geq q - 1$. This indicates that, for the Cauchy problem (1.1) with $\varphi \in H_q^{n/q}$, the critical growth of the nonlinearity at infinity should have the same rate as the case $\varphi \in \exp L^{q/(q-1)}$, that is, $f(u) \sim e^{u^{q/(q-1)}}$.

3 Preliminaries

In this section we recall some properties of the fundamental solution of the fractional diffusion equation, and give preliminary estimates. Furthermore we prove the boundedness of the nonlinear term $f(u)$ under the smallness assumption for the solution u .

Let $\theta \in (0, 2]$ and $G_\theta(t, x)$ be the fundamental solutions of the linear diffusion equations

$$\partial_t u + \mathcal{L}_\theta u = 0, \quad t > 0, \quad x \in \mathbb{R}^n, \quad (3.1)$$

where \mathcal{L}_θ is given in (2.13). It is well known that G_θ satisfies the following (see, i.e., [11]):

(i) $\int_{\mathbb{R}^n} G_\theta(t, x) dx = 1$ for any $t > 0$;

(ii) For any $j \in \mathbb{N}_0$,

$$|\nabla^j G_\theta(t, x)| \leq C t^{-\frac{n+j}{\theta}} \left(1 + t^{-1/\theta} |x|\right)^{-(n+j)}, \quad t > 0, \quad x \in \mathbb{R}^n.$$

Furthermore, for any $1 \leq r \leq \infty$,

$$\sup_{t>0} t^{\frac{n}{\theta}(1-\frac{1}{r})} \|G_\theta(t)\|_{L^r} < \infty;$$

(iii) For any $0 < s \leq t$,

$$G_\theta(t, x) = \int_{\mathbb{R}^n} G_\theta(t-s, x-y) G_\theta(s, y) dy. \quad (3.2)$$

For any $\varphi \in L^r$ ($1 \leq r \leq \infty$), we put

$$e^{-t\mathcal{L}_\theta}\varphi(x) := \int_{\mathbb{R}^n} G_\theta(t, x-y)\varphi(y) dy, \quad t > 0, \quad x \in \mathbb{R}^n,$$

which is a solution of (3.1) with the initial data φ , and it follows from (3.2) that

$$e^{-t\mathcal{L}_\theta}\varphi(x) = e^{-(t-s)\mathcal{L}_\theta}[e^{-s\mathcal{L}_\theta}\varphi](x), \quad t \geq s > 0, \quad x \in \mathbb{R}^n.$$

Combining property (ii) with the Young inequality and [11, 12], we have:

(G₁) There exists a constant c_θ , which depends only on n and θ , such that

$$\|e^{-t\mathcal{L}_\theta}\varphi\|_r \leq c_\theta t^{-\frac{n}{\theta}(\frac{1}{q}-\frac{1}{r})}\|\varphi\|_{L^q}, \quad \|e^{-t\mathcal{L}_\theta}\varphi\|_{L^q} \leq \|\varphi\|_{L^q}, \quad t > 0, \quad (3.3)$$

for any $\varphi \in L^q$ and $1 \leq q \leq r \leq \infty$;

(G₂) Let $\varphi \in L^1$ be such that

$$\int_{\mathbb{R}^n} \varphi(x) dx = 0.$$

Then

$$\lim_{t \rightarrow \infty} \|e^{-t\mathcal{L}_\theta}\varphi\|_{L^1} = 0.$$

We recall now the definition and the main properties of the Orlicz space $\exp L^r$ for $r > 1$.

Definition 3.1 Let $r > 1$. We define the Orlicz space $\exp L^r$ as

$$\exp L^r := \left\{ u \in L^1_{loc}(\mathbb{R}^n), \exists \lambda > 0 : \int_{\mathbb{R}^n} \left(\exp \left(\frac{|u(x)|}{\lambda} \right)^r - 1 \right) dx < \infty \right\},$$

where the norm is given by the Luxemburg type

$$\|u\|_{\exp L^r} := \inf \left\{ \lambda > 0 \text{ such that } \int_{\mathbb{R}^n} \left(\exp \left(\frac{|u(x)|}{\lambda} \right)^r - 1 \right) dx \leq 1 \right\}.$$

The space $\exp L^r$ endowed with the norm $\|u\|_{\exp L^r}$ is a Banach space. Moreover $\exp L^r \hookrightarrow L^q$ for any $q \in [r, \infty)$. An example of a function that is not bounded and that belongs to $\exp L^r$ is

$$u(x) = \begin{cases} (-\log|x|)^{1/r} & |x| < 1, \\ 0 & |x| \geq 1. \end{cases}$$

We stress also that C_0^∞ is not dense in $\exp L^r$. Finally the space $\exp L^r$ admits as predual the Orlicz space defined by the complementary function of $A(t) = e^{t^r} - 1$, denoted by $\tilde{A}(t)$. This complementary function is in particular a convex function such that $\tilde{A}(t) \sim t^2$ as $t \rightarrow 0$ and $\tilde{A}(t) \sim t \log^{1/r} t$ as $t \rightarrow \infty$. For more information about the Orlicz spaces, we refer to [1, 17]. Furthermore the following estimates hold.

Lemma 3.1 *Let $r > 1$. Then, for any $p \in [1, r]$, there exists a positive constant C_θ , which depends only on n and θ , such that*

$$\begin{aligned} \|e^{-t\mathcal{L}_\theta}\varphi\|_{\text{exp}L^r} &\leq \|\varphi\|_{\text{exp}L^r}, \quad t > 0, \\ \|e^{-t\mathcal{L}_\theta}\varphi\|_{\text{exp}L^r} &\leq C_\theta t^{-\frac{n}{\theta p}} \left(\log\left(t^{\frac{n}{\theta}} + 1\right)\right)^{-\frac{1}{r}} \|\varphi\|_{L^p}, \quad t > 0. \end{aligned}$$

A proof of this lemma is based on the following basic estimates and, for the case $r = 2$, can be found in [8, 9].

Lemma 3.2 *Let $r > 1$. Then it holds that*

$$\|\psi\|_{L^p} \leq \left[\Gamma\left(\frac{p}{r} + 1\right)\right]^{\frac{1}{p}} \|\psi\|_{\text{exp}L^r} \quad (3.4)$$

for any $p \in [r, \infty)$, where Γ is the Gamma function

$$\Gamma(q) := \int_0^\infty \xi^{q-1} e^{-\xi} d\xi, \quad q > 0.$$

A proof of this lemma can be found in [19].

Lemma 3.3 *For any $p \geq 1$ and $r \geq 1$, there exists a positive constant C , which is independent of p and r , such that*

$$\Gamma(rp + 1)^{\frac{1}{p}} \leq C\Gamma(r + 1)p^r. \quad (3.5)$$

Proof. This is a consequence of Stirling's formula. Indeed, for a fixed positive constant $C > 1$, there exists r_0 large enough such that, for any $r \geq r_0$ and $p \geq 1$,

$$\frac{\Gamma(rp + 1)^{1/p}}{\Gamma(r + 1)} \leq C \frac{(rp)^r e^{-r} (2\pi rp)^{1/(2p)}}{r^r e^{-r} (2\pi r)^{1/2}} \leq Cp^r.$$

For $1 \leq r \leq r_0$, we consider first the case of large values of p , namely $p \geq p_0$, and we obtain an estimate similar to the previous one. Finally we observe that for $1 \leq r \leq r_0$ and $1 \leq p \leq p_0$ the quotient is bounded by a constant. \square

At the end of this section we give the following estimate on the nonlinearity $f(u)$, which is crucial throughout this paper.

Lemma 3.4 *Let $n \geq 1$ and $M > 0$. Suppose that the function $u \in L^\infty(0, \infty; \text{exp}L^2)$ satisfies the condition*

$$\sup_{t>0} \|u(t)\|_{\text{exp}L^2} \leq M.$$

Let f be the function defined as in (1.6). Furthermore, let p_1 be the constant given in (2.8), namely $p_1 = \max(1, (2n)/(4 + n))$. Then, for all $p \in [p_1, \infty)$, there exists $\varepsilon = \varepsilon(p) > 0$ such that, if $M < \varepsilon$, then

$$\sup_{t>0} \|f(u(t))\|_{L^p} \leq 2Cp^3 M^{1+\frac{4}{n}}, \quad (3.6)$$

where C is independent of p , n and M .

Proof. For any $k \in \mathbb{N}_0$, we put

$$\ell_k := 2k + 1 + \frac{4}{n}. \quad (3.7)$$

Then, since it holds from $p \geq p_1$ with (2.8) that

$$\ell_k p \geq \left(1 + \frac{4}{n}\right) \frac{2n}{n+4} = 2$$

for any $k \in \mathbb{N}_0$, by (1.6) and (3.4) we have

$$\begin{aligned} \|f(u(t))\|_{L^p} &\leq \sum_{k=0}^{\infty} \frac{\|u^{\ell_k}(t)\|_{L^p}}{k!} \\ &= \sum_{k=0}^{\infty} \frac{\|u(t)\|_{L^{\ell_k p}}^{\ell_k}}{k!} \leq \sum_{k=0}^{\infty} \frac{\left(\Gamma\left(\frac{\ell_k p}{2} + 1\right)^{\frac{1}{\ell_k p}} \|u(t)\|_{\text{exp}L^2}\right)^{\ell_k}}{k!} \end{aligned} \quad (3.8)$$

for all $t > 0$. By (3.5) with the monotonicity property of the Gamma function we see that

$$\begin{aligned} \Gamma\left(\frac{\ell_k p}{2} + 1\right)^{\frac{1}{p}} &\leq C \Gamma\left(\frac{\ell_k}{2} + 1\right) p^{\frac{\ell_k}{2}} \\ &= C \Gamma\left(k + \frac{3}{2} + \frac{2}{n}\right) p^{\frac{\ell_k}{2}} \\ &\leq C \Gamma(k+4) p^{\frac{\ell_k}{2}} = C(k+3)! p^{\frac{\ell_k}{2}}. \end{aligned}$$

This together with the assumption on u and (3.8) implies that

$$\begin{aligned} \|f(u(t))\|_{L^p} &\leq C \sum_{k=0}^{\infty} \frac{(k+3)!}{k!} \left(p \|u(t)\|_{\text{exp}L^2}^2\right)^{\frac{\ell_k}{2}} \\ &\leq C \sum_{k=0}^{\infty} (k+1)(k+2)(k+3) (pM^2)^{k+\frac{1}{2}+\frac{2}{n}}, \quad t > 0. \end{aligned}$$

Now for $M < \varepsilon(p)$ small enough we get

$$\sup_{t>0} \|f(u(t))\|_{L^p} \leq C \frac{(pM^2)^{\frac{1}{2}+\frac{2}{n}}}{(1-pM^2)^4} \leq 2C(pM^2)^{\frac{1}{2}+\frac{2}{n}} \leq 2Cp^3 M^{1+\frac{4}{n}}.$$

This implies (3.6), and the proof of Lemma 3.4 is complete. \square

4 Equivalence and uniqueness

In this section we prove the equivalence between small weak and small mild solutions of (1.1). Furthermore, we show that the small solution is unique in any dimension $n \in \mathbb{N}$.

We first prove Proposition 2.1.

Proof of Proposition 2.1. Let us first remark that, for ε small enough, by (1.1) we can apply Lemma 3.4, and we see that $f(u) \in L^\infty(0, \infty; L^{p_1})$ where p_1 is the constant given in (2.8).

Suppose that u is a weak solution of the Cauchy problem (1.1). Then, for $0 < s < t$, it holds

$$e^{(t-s)\Delta} f(u(s)) = e^{(t-s)\Delta} (\partial_s u - \Delta u) = \partial_s \left(e^{(t-s)\Delta} u(s) \right) \quad \text{in } \mathcal{D}'.$$

Integrating on (τ, t) , we obtain

$$u(t) = e^{(t-\tau)\Delta} u(\tau) + \int_\tau^t e^{(t-s)\Delta} f(u(s)) ds$$

for all $t > \tau > 0$. Let $\tau \rightarrow 0$. Since $f(u) \in L^\infty(0, T; L^{p_1})$, we get

$$\int_\tau^t e^{(t-s)\Delta} f(u(s)) ds \rightarrow \int_0^t e^{(t-s)\Delta} f(u(s)) ds \quad \text{in } L^{p_1}.$$

Moreover, it follows from $u(\tau) \xrightarrow{*} \varphi$ in $\exp L^2$ that

$$e^{(t-\tau)\Delta} u(\tau) \xrightarrow{*} e^{t\Delta} \varphi,$$

and the same limit holds in \mathcal{D}' . So we obtain

$$u(t) = e^{t\Delta} \varphi + \int_0^t e^{(t-s)\Delta} f(u(s)) ds \quad \text{in } \mathcal{D}'.$$

Moreover $e^{t\Delta} \varphi \in C((0, \infty); \exp L^2)$ and by Lemma 3.4 the integral term belongs to $C([0, \infty), \exp L^2)$. This completes the proof that statement (i) implies statement (ii).

Suppose now that $u \in L^\infty(0, T; \exp L^2)$ is a mild solution of (1.1). We know that

$$\partial_t e^{t\Delta} \varphi = \Delta e^{t\Delta} \varphi \quad \text{in } \mathcal{D}'.$$

Let us consider $\eta \in \mathcal{D}((0, T) \times \mathbb{R}^n)$. Of course, there exists $\tilde{T} \in (0, T)$ such that $\eta(t, x) = 0$ for $t \in [\tilde{T}, T)$. By a direct computation we are going to prove that

$$\begin{aligned} & \langle \partial_t \int_0^t e^{(t-s)\Delta} f(u(s)) ds, \eta(t) \rangle_{L^2(L^2)} \\ &= \langle f(u), \eta \rangle_{L^2(L^2)} + \langle \Delta \int_0^t e^{(t-s)\Delta} f(u(s)) ds, \eta(t) \rangle_{L^2(L^2)}. \end{aligned}$$

Indeed,

$$\begin{aligned}
& \langle \partial_t \int_0^t e^{(t-s)\Delta} f(u(s)) ds, \eta(t) \rangle_{L^2(L^2)} \\
&= - \langle \int_0^t e^{(t-s)\Delta} f(u(s)) ds, \partial_t \eta(t) \rangle_{L^2(L^2)} \\
&= - \int_0^{\tilde{T}} \langle \int_0^t e^{(t-s)\Delta} f(u(s)) ds, \partial_t \eta(t) \rangle_{L^2} dt \\
&= - \int_0^{\tilde{T}} \int_0^t \langle e^{(t-s)\Delta} f(u(s)), \partial_t \eta(t) \rangle_{L^2} ds dt \\
&= - \int_0^{\tilde{T}} \int_0^t \langle f(u(s)), e^{(t-s)\Delta} \partial_t \eta(t) \rangle_{L^2} ds dt \\
&= \int_0^{\tilde{T}} \int_0^t \langle f(u(s)), -\partial_t (e^{(t-s)\Delta} \eta(t)) + \Delta e^{(t-s)\Delta} \eta(t) \rangle_{L^2} ds dt \\
&= \int_0^{\tilde{T}} \int_0^t -\partial_t \langle f(u(s)), e^{(t-s)\Delta} \eta(t) \rangle_{L^2} ds dt + \int_0^{\tilde{T}} \int_0^t \langle f(u(s)), \Delta e^{(t-s)\Delta} \eta(t) \rangle_{L^2} ds dt \\
&= \int_0^{\tilde{T}} \int_s^{\tilde{T}} -\partial_t \langle f(u(s)), e^{(t-s)\Delta} \eta(t) \rangle_{L^2} dt ds + \int_0^{\tilde{T}} \langle \int_0^t \Delta e^{(t-s)\Delta} f(u(s)) ds, \eta(t) \rangle_{L^2} dt \\
&= \int_0^{\tilde{T}} \langle f(u(s)), \eta(s) \rangle_{L^2} ds + \langle \Delta \int_0^t e^{(t-s)\Delta} f(u(s)) ds, \eta(t) \rangle_{L^2(L^2)} \\
&= \langle f(u), \eta \rangle_{L^2(L^2)} + \langle \Delta \int_0^t e^{(t-s)\Delta} f(u(s)) ds, \eta(t) \rangle_{L^2(L^2)}.
\end{aligned}$$

Therefore we obtain

$$\partial_t u = \Delta u + f(u) \quad \text{in } \mathcal{D}'.$$

This yields that statement (ii) implies statement (i), and the proof of Theorem 2.1 is complete. \square

Next we show the uniqueness of the small solution of (1.1).

Proof of Proposition 2.2. Let us suppose that u, v are two solutions of the Cauchy problem (1.1). Furthermore, since they converge to φ weak* in $\text{exp}L^2$, we can continue them in $t = 0$ as φ . Moreover, by Lemma 3.4 we have

$$\|u(t) - v(t)\|_{\text{exp}L^2} \leq \|u(t) - e^{t\Delta} \varphi\|_{\text{exp}L^2} + \|v(t) - e^{t\Delta} \varphi\|_{\text{exp}L^2} \rightarrow 0 \quad \text{as } t \rightarrow +0.$$

as $t \rightarrow +0$. Let

$$H(t) := \|u(t) - v(t)\|_{\text{exp}L^2}, \quad t \geq 0, \tag{4.1}$$

so that $H(0) = 0$. Furthermore, put

$$t_0 := \sup \{t \in [0, \min(T, T')]\} \text{ such that } H(s) = 0 \text{ for every } s \in [0, t].$$

By (4.1) we have $t_0 \geq 0$. By contradiction we assume $t_0 < \min(T, T')$. Since $H(t)$ is continuous in time we have $H(t_0) = 0$. Let us denote $\tilde{u}(t) = u(t + t_0)$ and $\tilde{v}(t) = v(t + t_0)$

so that \tilde{u} and \tilde{v} satisfy equation (1.1) on $(0, \infty)$ and by $H(t_0) = 0$ it follows $\tilde{u}(0) = \tilde{v}(0)$. We will prove that there exists a positive time \tilde{t} such that

$$\sup_{0 < t < \tilde{t}} \|\tilde{u}(t) - \tilde{v}(t)\|_{\text{exp}L^2} \leq C(\tilde{t}) \sup_{0 < t < \tilde{t}} \|\tilde{u}(t) - \tilde{v}(t)\|_{\text{exp}L^2} \quad (4.2)$$

for a constant $C(\tilde{t}) < 1$, and so $\|\tilde{u}(t) - \tilde{v}(t)\|_{\text{exp}L^2} = 0$ for any $t \in [0, \tilde{t}]$. Therefore $u(t + t_0) = v(t + t_0)$ for any $t \in [0, \tilde{t}]$ in contradiction with the definition of t_0 . In order to establish inequality (4.2) we control both the L^2 norm and the L^∞ norm of the difference of the two solutions. By (1.6) we see that there exist positive constants C and λ such that

$$|f(x) - f(y)| \leq C|x - y| \left(e^{\lambda x^2} + e^{\lambda y^2} \right), \quad x, y \in \mathbb{R}.$$

Then, for $p, q > 2$ such that $1/p + 1/q = 1/2$, we have

$$\begin{aligned} \|\tilde{u}(t) - \tilde{v}(t)\|_{L^2} &\leq C \int_0^t \left\| |\tilde{u}(s) - \tilde{v}(s)| (e^{\lambda \tilde{u}^2(s)} + e^{\lambda \tilde{v}^2(s)}) \right\|_{L^2} ds \\ &\leq C \int_0^t \|\tilde{u}(s) - \tilde{v}(s)\|_{L^2} ds \\ &\quad + C \int_0^t \left(\|\tilde{u}(s) - \tilde{v}(s)\|_{L^q} \left\| (e^{\lambda \tilde{u}^2(s)} - 1) + (e^{\lambda \tilde{v}^2(s)} - 1) \right\|_{L^p} \right) ds \\ &\leq Ct \sup_{0 < s < t} \|\tilde{u}(s) - \tilde{v}(s)\|_{\text{exp}L^2} \\ &\quad + C \sup_{0 < s < t} \|\tilde{u}(s) - \tilde{v}(s)\|_{\text{exp}L^2} \int_0^t \left\| (e^{\lambda \tilde{u}^2(s)} - 1) + (e^{\lambda \tilde{v}^2(s)} - 1) \right\|_{L^p} ds \end{aligned}$$

for all $t > 0$. Moreover, if $\sup_{t > 0} \|\tilde{u}(t)\|_{\text{exp}L^2}$ and $\sup_{t > 0} \|\tilde{v}(t)\|_{\text{exp}L^2}$ are small enough, then, by the same proof as Lemma 3.4 the term in the integral is uniformly bounded in time. Indeed,

$$\begin{aligned} \sup_{0 < s < \infty} \left\| (e^{\lambda \tilde{u}^2(s)} - 1) + (e^{\lambda \tilde{v}^2(s)} - 1) \right\|_{L^p} \\ \leq C(\tilde{u}, \tilde{v}) < \infty. \end{aligned} \quad (4.3)$$

Therefore, we obtain

$$\sup_{0 < s < t} \|\tilde{u}(s) - \tilde{v}(s)\|_{L^2} \leq C(\tilde{u}, \tilde{v}) t \sup_{0 < s < t} \|\tilde{u}(s) - \tilde{v}(s)\|_{\text{exp}L^2}. \quad (4.4)$$

In a similar way, for $r > 2$ such that $n/(2r) < 1$, we get

$$\begin{aligned} \|\tilde{u}(t) - \tilde{v}(t)\|_{L^\infty} &\leq C \int_0^t (t-s)^{-\frac{n}{2r}} \left\| |\tilde{u}(s) - \tilde{v}(s)| \left(e^{\lambda \tilde{u}^2(s)} + e^{\lambda \tilde{v}^2(s)} \right) \right\|_{L^r} ds \\ &\leq C \int_0^t (t-s)^{-\frac{n}{2r}} \|\tilde{u}(s) - \tilde{v}(s)\|_{L^r} ds \\ &\quad + C \int_0^t (t-s)^{-\frac{n}{2r}} \|\tilde{u}(s) - \tilde{v}(s)\|_{L^q} \left\| e^{\lambda \tilde{u}^2(s)} - 1 + e^{\lambda \tilde{v}^2(s)} - 1 \right\|_{L^p} ds \end{aligned}$$

for some \bar{q}, \bar{p} such that $1/\bar{q} + 1/\bar{p} = 1/r$. Since $\bar{p} \geq r > 2$, we can apply an estimate similar to (4.3), and obtain that

$$\sup_{0 < s < t} \|\tilde{u}(s) - \tilde{v}(s)\|_{L^\infty} \leq C(\tilde{u}, \tilde{v}) t^{1-\frac{n}{2r}} \sup_{0 < s < t} \|\tilde{u}(s) - \tilde{v}(s)\|_{\exp L^2}. \quad (4.5)$$

Therefore the two inequalities (4.4) and (4.5) imply

$$\sup_{0 < s < t} \|\tilde{u}(s) - \tilde{v}(s)\|_{\exp L^2} \leq C(\tilde{u}, \tilde{v})(t^{1-\frac{n}{2r}} + t) \sup_{0 < s < t} \|\tilde{u}(s) - \tilde{v}(s)\|_{\exp L^2},$$

and, for t small enough, we obtain the desired estimate. \square

5 Singular initial data

In order to prove Theorem 2.1 the principal idea is the following. Even if the initial data φ satisfy $\varphi \in \exp L^2$ and so $\varphi \in L^p$ for $p \in [2, \infty)$ with a norm blowing up with p , the solution is indeed in $L^2 \cap L^\infty$ for all $t > 0$. Moreover, for each fixed $t_0 > 0$, the norm $\|u(t_0)\|_\infty$ is arbitrarily small provided the initial data are sufficiently small in $\exp L^2$. So, thanks to the uniqueness result (Proposition 2.2), it is possible to consider the solution $u(t)$ for $t \geq t_0$ as the limit of a recursive procedure building up a solution starting with initial data belonging to all L^p with $p \in [2, \infty]$ and with norm uniformly bounded.

Let us prove now (2.3) for small time.

Lemma 5.1 *Let $n \geq 1$. There is $\varepsilon = \varepsilon(n) > 0$ such that, if $\|\varphi\|_{\exp L^2} < \varepsilon$, then the unique solution u of the Cauchy problem (1.1) satisfying (1.8) and*

$$\|u(t)\|_{L^q} \leq Ct^{-\frac{n}{2}\left(\frac{1}{2}-\frac{1}{q}\right)} \|\varphi\|_{\exp L^2}, \quad 0 < t \leq 1, \quad (5.1)$$

for any $q \in [2, \infty]$, where $C > 0$ depending only on n .

Proof. Let us consider

$$u(t) = e^{t\Delta}\varphi + \int_0^t e^{(t-s)\Delta} f(u(s)) ds := e^{t\Delta}\varphi + F(u(t))$$

and let us denote as in Lemma 3.4

$$M = 2\|\varphi\|_{\exp L^2}.$$

As for the linear part, for any $q \in [2, \infty]$, by (3.3) and (3.4) we have

$$\|e^{t\Delta}\varphi\|_{L^q} \leq c_2 t^{-\frac{n}{2}\left(\frac{1}{2}-\frac{1}{q}\right)} \|\varphi\|_{L^2} \leq c_2 t^{-\frac{n}{2}\left(\frac{1}{2}-\frac{1}{q}\right)} \|\varphi\|_{\exp L^2}, \quad t > 0.$$

Let us consider now the the nonlinear part $F(u)$. By choosing $r \geq p_1$ such that $2r > n$ (and so r depends only on the dimension n), we get

$$\|F(u(t))\|_{L^\infty} \leq c_2 \int_0^t (t-s)^{-\frac{n}{2r}} \|f(u(s))\|_{L^r} ds \leq Ct^{1-\frac{n}{2r}} \sup_{t>0} \|f(u(t))\|_{L^r}, \quad t > 0.$$

Now, by Lemma 3.4 we get for M small enough depending on r (and so on n only)

$$Ct^{1-\frac{n}{2r}} \sup_{t>0} \|f(u(t))\|_{L^r} \leq Cr^3 t^{1-\frac{n}{2r}} M^{1+\frac{4}{n}}.$$

Now, for $t \in (0, 1]$ and M small enough, depending again only on n , we get

$$\sup_{0<t\leq 1} \|F(u(t))\|_{L^\infty} \leq C\|\varphi\|_{\exp L^2}.$$

So we get

$$\|u(t)\|_{L^\infty} \leq c_2 t^{-\frac{n}{4}} \|\varphi\|_{\exp L^2} + C\|\varphi\|_{\exp L^2} \leq C t^{-\frac{n}{4}} \|\varphi\|_{\exp L^2}, \quad t \in (0, 1]. \quad (5.2)$$

On the other hand, by the embedding $\exp L^2 \subset L^2$ (3.4) we get

$$\|u(t)\|_{L^2} \leq \Gamma(2)^{\frac{1}{2}} \|u(t)\|_{\exp L^2}, \quad t > 0.$$

This together with (5.2) implies (5.1), and the proof of Lemma 5.1 is complete. \square

We have now to prove (2.3) for large times $t > 1$. Let us put

$$v(t, x) := u(t + 1, x)$$

so that

$$\begin{cases} \partial_t v = \Delta v + f(v), & t > 0, \quad x \in \mathbb{R}^n \\ v(0, x) = u(1, x) \in L^q, & x \in \mathbb{R}^n, \quad q \in [2, \infty], \end{cases}$$

and by Lemma 5.1 for all $q \in [2, \infty]$ we have $\|u(1)\|_{L^q} \leq C\|\varphi\|_{\exp L^2} \leq \varepsilon$ small enough. Let us consider now the following recursive sequence $\{v_j\}_{j \in \mathbb{N}}$

$$\begin{cases} v_0(t) = e^{t\Delta} v(0) \\ v_{j+1}(t) = v_0(t) + \int_0^t e^{(t-s)\Delta} f(v_j(s)) ds, \quad j \geq 0. \end{cases} \quad (5.3)$$

Since $0 \leq v_j(t, x) \leq v_{j+1}(t, x) \leq v(t, x)$ for $x \in \mathbb{R}^n$ and $t > 0$, we get by uniqueness

$$\lim_{j \rightarrow \infty} v_j(t, x) = v(t, x) = u(t + 1, x) \quad (5.4)$$

the limit being considered pointwise. We prove now (2.3) for large times through the recursive sequence (5.3). We begin by a Lemma which describes how boundedness and decreasing in time propagate on the nonlinear term.

Lemma 5.2 *Let $n \geq 1$. Let $M = 2\|\varphi\|_{\exp L^2}$ and $v(x, t)$ a function satisfying*

$$\sup_{s>0} (1+s)^{\frac{n}{2}} \left(\frac{1}{2}-\frac{1}{q}\right) \|v(s)\|_{L^q} \leq CM, \quad q \in [2, \infty], \quad (5.5)$$

where C independent of q . Then, there exists $\varepsilon > 0$ such that, if $M < \varepsilon$, then, for any $r \in [p_1, \infty]$,

$$\sup_{s>0} (1+s)^{\frac{n}{2}} \left(\frac{1}{2}-\frac{1}{r}\right)^{+1} \|f(v(s))\|_{L^r} \leq 2(CM)^{1+\frac{4}{n}} \quad (5.6)$$

where f and p_1 are defined as in (1.6) and (2.8), respectively.

Proof. Let $k \in \mathbb{N}_0$ and ℓ_k be the constant given in (3.7). Then, for any $r \in [p_1, \infty]$, by (1.6) and (5.5) we have

$$\begin{aligned}
\|f(v(s))\|_{L^r} &\leq \sum_{k=0}^{\infty} \frac{1}{k!} \|v(s)\|_{L^{\ell_k r}}^{\ell_k} \\
&\leq \sum_{k=0}^{\infty} \frac{1}{k!} \left(C(1+s)^{-\frac{n}{2} \left(\frac{1}{2} - \frac{1}{\ell_k r} \right)} M \right)^{\ell_k} \\
&\leq (CM)^{1+\frac{4}{n}} (1+s)^{\frac{n}{2r} - \frac{n}{4} \left(1 + \frac{4}{n} \right)} \sum_{k=0}^{\infty} \frac{1}{k!} \left(C(1+s)^{-\frac{n}{4}} M \right)^{2k} \\
&\leq (CM)^{1+\frac{4}{n}} (1+s)^{-\frac{n}{2} \left(\frac{1}{2} - \frac{1}{r} \right) - 1} \sum_{k=0}^{\infty} \frac{1}{k!} (CM)^{2k}
\end{aligned} \tag{5.7}$$

for all $s > 0$. We can take a sufficiently small ε , which is independent of r , so that, for $M \leq \varepsilon$, it holds that

$$\sum_{k=0}^{\infty} \frac{1}{k!} (CM)^{2k} = e^{(CM)^2} \leq 2.$$

This together with (5.7) implies (5.6). Thus Lemma 5.2 follows. \square

Now we are in position to prove Theorem 2.1.

Proof of Theorem 2.1. By Lemma 5.1 we have (2.3) for $0 < t \leq 1$. So it suffices to prove (2.3) for $t > 1$. Let $j \in \mathbb{N}_0$ and $v_j(t, x)$ be the function given in (5.3). Since $v_0(t) = e^{t\Delta} v(0) = e^{t\Delta} u(1)$, for any $q \in [2, \infty]$, we have by (3.3)

$$\|v_0(t)\|_{L^q} \leq c_2 t^{-\frac{n}{2} \left(\frac{1}{2} - \frac{1}{q} \right)} \|v(0)\|_{L^2}, \quad t > 0,$$

and also

$$\|v_0(t)\|_{L^q} \leq \|v(0)\|_{L^q}, \quad t > 0.$$

On the other hand, by Lemma 5.1 we have

$$\|v(0)\|_{L^q} = \|u(1)\|_{L^q} \leq C \|\varphi\|_{\exp L^2}.$$

Therefore, for any $q \in [2, \infty]$, there exists a positive constant c_* depending only on n such that

$$\|v_0(t)\|_{L^q} \leq c_* (1+t)^{-\frac{n}{2} \left(\frac{1}{2} - \frac{1}{q} \right)} \|\varphi\|_{\exp L^2}. \tag{5.8}$$

Let $m \in \mathbb{N}_0$, and put

$$\tilde{M} := c_* \|\varphi\|_{\exp L^2}.$$

Let us assume that, for $j = m$,

$$\|v_m(t)\|_{L^q} \leq 2\tilde{M} (1+t)^{-\frac{n}{2} \left(\frac{1}{2} - \frac{1}{q} \right)}, \quad t > 0, \tag{5.9}$$

and let us prove it for $j = m + 1$. By (5.3) it suffices to consider the nonlinear term. For any $q \in [2, \infty]$, we put

$$\begin{aligned} & \left\| \int_0^t e^{(t-s)\Delta} f(v_m(s)) ds \right\|_{L^q} \\ & \leq \int_0^{t/2} \left\| e^{(t-s)\Delta} f(v_m(s)) \right\|_{L^q} ds + \int_{t/2}^t \left\| e^{(t-s)\Delta} f(v_m(s)) \right\|_{L^q} ds \\ & =: J_1(t) + J_2(t), \quad t > 0. \end{aligned}$$

For the term $J_1(t)$, by (3.3) and (5.6) we obtain

$$\begin{aligned} J_1(t) & \leq \int_0^{t/2} \|f(v_m(s))\|_{L^q} ds \\ & \leq 2C\tilde{M}^{1+\frac{4}{n}} \int_0^1 (1+s)^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{q})-1} ds \leq 2C\tilde{M}^{1+\frac{4}{n}} \end{aligned} \tag{5.10}$$

for all $0 < t \leq 2$. Moreover, since $1 \leq p_1 < 2$, by (3.3) and (5.6) we have

$$\begin{aligned} J_1(t) & \leq c_2 \int_0^{t/2} (t-s)^{-\frac{n}{2}(\frac{1}{p_1}-\frac{1}{q})} \|f(v_m(s))\|_{L^{p_1}} ds \\ & \leq C\tilde{M}^{1+\frac{4}{n}} t^{-\frac{n}{2}(\frac{1}{p_1}-\frac{1}{q})} \int_0^{\frac{t}{2}} (1+s)^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{p_1})-1} ds \\ & \leq C\tilde{M}^{1+\frac{4}{n}} t^{-\frac{n}{2}(\frac{1}{p_1}-\frac{1}{q})} (1+t)^{\frac{n}{2}(\frac{1}{p_1}-\frac{1}{2})} \leq C\tilde{M}^{1+\frac{4}{n}} t^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{q})} \end{aligned}$$

for all $t \geq 1$. This together with (5.10) implies that

$$J_1(t) \leq D_1 \tilde{M}^{1+\frac{4}{n}} (1+t)^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{q})}, \quad t > 0,$$

where D_1 is a positive constant independent of m and t . On the other hand, for the term $J_2(t)$, exploiting (3.3) and (5.6) again, we see that

$$\begin{aligned} J_2(t) & \leq \int_{t/2}^t \|f(v_m(s))\|_{L^q} ds \leq 2C\tilde{M}^{1+\frac{4}{n}} \int_{t/2}^t (1+s)^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{q})-1} ds \\ & \leq D_2 \tilde{M}^{1+\frac{4}{n}} (1+t)^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{q})}, \quad t > 0, \end{aligned}$$

where D_2 is a positive constant which independent of m and t . We can now assume

$$(D_1 + D_2) \tilde{M}^{\frac{4}{n}} = (D_1 + D_2) (c_* \|\varphi\|_{\text{exp}L^2})^{\frac{4}{n}} < 1$$

in order to get

$$\left\| \int_0^t e^{(t-s)\Delta} f(v_m(s)) ds \right\|_{L^q} \leq J_1(t) + J_2(t) \leq \tilde{M} (1+t)^{\frac{n}{2}(\frac{1}{2}-\frac{1}{q})}, \quad t > 0. \tag{5.11}$$

By (5.3), (5.8) and (5.11) we get the estimate (5.9) on v_{m+1} . Therefore we have the estimate (5.9) for all $m \in \mathbb{N}_0$. Passing to the limit with (5.4) we see that

$$\|u(t+1)\|_{L^q} = \|v(t)\|_{L^q} \leq 2c_*(1+t)^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{q})} \|\varphi\|_{\text{exp}L^2}, \quad t > 0.$$

This implies (2.3) for $t > 1$, and the proof of Theorem 2.1 is complete. \square

6 Regular initial data

We are going to prove now Theorem 2.2. For the estimate (2.6) for small times, we are going to exploit again Lemma 3.4 but this time, in order not to assume smallness on $\|\varphi\|_{L^p}$, we introduce a time T_* , to be suitably chosen, and we will split the estimate (2.6) into $0 < t \leq 2T_*$ and $t > 2T_*$. The same splitting would have worked also in the singular case without leading however to any significative improvement. Let

$$M := \|\varphi\|_{\text{exp}L^2}.$$

By Proposition 1.1, it is not restrictive to assume $M < 1$. For $\|\varphi\|_{p_*} > 0$ we have

$$M \leq 2\|\varphi\|_{\text{exp}L^2 \cap L^{p_*}} \leq 2 \max\{2M, 2\|\varphi\|_{L^{p_*}}\} \leq 4 \max\{1, \|\varphi\|_{L^{p_*}}\}.$$

Let now

$$K := 4 \max\{1, \|\varphi\|_{L^{p_*}}\} \quad (6.1)$$

and so

$$M \leq K. \quad (6.2)$$

Then we begin by a Lemma analogous to Lemma 5.2.

Lemma 6.1 *Let $n \geq 1$ and $p \in [1, 2)$. Furthermore let p_* be the constant given in (2.4). Suppose that $v(t, x)$ is a function satisfying*

$$\sup_{s>0} s^{\frac{n}{2}(\frac{1}{p_*} - \frac{1}{q})} \|v(s)\|_{L^q} \leq CK$$

for any $q \in [p_*, \infty]$, where C is independent of q . Then, there exists a sufficiently large constant $T_1 = T_1(K, p_*) > 1$ such that, for any $r \in [p_2, \infty]$,

$$\sup_{s>T_1} s^{\frac{n}{2}(\frac{1}{p_*} - \frac{1}{r}) + \frac{2}{p_*}} \|f(v(s))\|_{L^r} \leq 2(CK)^{1 + \frac{4}{n}}, \quad (6.3)$$

where f is defined as in (1.6) and

$$p_2 = \max\left\{\frac{p_* n}{n+4}, 1\right\}. \quad (6.4)$$

Remark 6.1 *Comparing with Lemma 5.2, since we are not assuming boundedness of $\|v(s)\|_{L^q}$ near $s = 0$, we can only obtain boundedness and decreasing of the nonlinear term for large times.*

Remark 6.2 *For $1 \leq n \leq 4$, since $p \in [1, 2)$ and $(2n)/(n+4) \leq 1$, we have $p_* = p$ and $p_2 = 1$. On the contrary for $n \geq 5$, then p_* might be strictly greater than p and p_2 might be strictly greater than 1.*

Proof. Let $\ell_k = 2k + 1 + 4/n$. Since

$$\ell_k p_2 \geq \left(1 + \frac{4}{n}\right) p_2 \geq p_*,$$

for any $r \in [p_2, \infty]$, we see that

$$\begin{aligned}
\|f(v(s))\|_{L^r} &\leq \sum_{k=0}^{\infty} \frac{1}{k!} \|v(s)\|_{L^{\ell_k r}}^{\ell_k} \\
&\leq \sum_{k=0}^{\infty} \frac{1}{k!} \left(C s^{-\frac{n}{2}(\frac{1}{p_*} - \frac{1}{\ell_k r})} K \right)^{\ell_k} \\
&\leq (CK)^{1+\frac{4}{n}} s^{\frac{n}{2r} - \frac{n}{2p_*}(1+\frac{4}{n})} \sum_{k=0}^{\infty} \frac{\left(C s^{-\frac{n}{2p_*}} K \right)^{2k}}{k!} \\
&\leq (CK)^{1+\frac{4}{n}} s^{-\frac{n}{2}(\frac{1}{p_*} - \frac{1}{r}) - \frac{2}{p_*}} \exp\left((CK)^2 s^{-\frac{n}{p_*}} \right)
\end{aligned} \tag{6.5}$$

for all $s > 0$. We can choose a sufficiently large constant $T_1 \geq 1$ such that, for all $s > T_1$, it holds that

$$\exp\left((CK)^2 s^{-\frac{n}{p_*}} \right) \leq 2.$$

It is enough to chose

$$T_1 \geq \left(\frac{(CK)^2}{\log 2} \right)^{\frac{p_*}{n}}. \tag{6.6}$$

This together with (6.5) implies (6.3). Thus Lemma 6.1 follows. \square

Let us prove now (2.6) for small times.

Lemma 6.2 *Let $n \geq 1$ and $M = \|\varphi\|_{\text{exp}L^2} < \varepsilon$ small enough so that existence and uniqueness theorems apply. Furthermore, let u be the unique solution of the Cauchy problem (1.1) satisfying (1.8). Suppose $\varphi \in L^p$ for $p \in [1, 2)$ and let p_* be the constant given in (2.4). Then, for any fixed $T_2 \geq 1$, there exists a sufficiently small constant $\varepsilon = \varepsilon(p_*, T_2) > 0$ such that, if $M < \varepsilon$, then the solution u satisfies*

$$\|u(t)\|_{L^q} \leq \tilde{C} t^{-\frac{n}{2}(\frac{1}{p_*} - \frac{1}{q})} K, \quad 0 < t \leq 2T_2 \tag{6.7}$$

for all $q \in [p_*, \infty]$ and for $\tilde{C} > 0$ depending on n , where K is the constant defined by (6.1).

Proof. Let us consider

$$u(t) = e^{t\Delta} \varphi + \int_0^t e^{(t-s)\Delta} f(u(s)) ds := e^{t\Delta} \varphi + F(u(t)).$$

As for the linear part, for any $q \in [p_*, \infty]$, by (3.3) and (6.1) we have

$$\|e^{t\Delta} \varphi\|_{L^q} \leq c_2 t^{-\frac{n}{2}(\frac{1}{p_*} - \frac{1}{q})} \|\varphi\|_{L^{p_*}} \leq c_2 t^{-\frac{n}{2}(\frac{1}{p_*} - \frac{1}{q})} K, \quad t > 0. \tag{6.8}$$

Let us consider now the nonlinear part $F(u)$. For $r \geq p_1$ (defined in (2.8)) such that $2r > n$ (and so r depends on n only), we have

$$\|F(u(t))\|_{L^\infty} \leq c_2 \int_0^t (t-s)^{-\frac{n}{2r}} \|f(u(s))\|_{L^r} ds \leq C t^{1-\frac{n}{2r}} \sup_{t>0} \|f(u(t))\|_{L^r}, \quad t > 0,$$

where $C > 0$ depends on n only. Now, by Lemma 3.4, for $M < 1$ small enough depending only on r , we get

$$Ct^{1-\frac{n}{2r}} \sup_{t>0} \|f(u(t))\|_{L^r} \leq Ct^{1-\frac{n}{2r}} r^3 M^{1+\frac{4}{n}} \leq CT^{1-\frac{n}{2r}} M^{1+\frac{4}{n}}, \quad t \leq 2T_2,$$

where $C > 0$ depends on n only. On the other hand, since it follows from $p \in [1, 2)$ with (2.8) that

$$1 - \frac{n}{2} \left(\frac{1}{p_1} - \frac{1}{p_*} \right) > 0,$$

again by Lemma 3.4, for $M < 1$ small enough depending only on p_1 , we have

$$\begin{aligned} \|F(u(t))\|_{L^{p_*}} &\leq c_2 \int_0^t (t-s)^{-\frac{n}{2}(\frac{1}{p_1}-\frac{1}{p_*})} \|f(u(s))\|_{L^{p_1}} ds \\ &\leq Cp_1^3 t^{1-\frac{n}{2}(\frac{1}{p_1}-\frac{1}{p_*})} M^{1+\frac{4}{n}} \leq CT_2^{1-\frac{n}{2}(\frac{1}{p_1}-\frac{1}{p_*})} M^{1+\frac{4}{n}}, \quad t \leq 2T_2, \end{aligned}$$

where $C > 0$ depends on n . If we choose

$$M^{\frac{4}{n}} \max(T_2^{1-\frac{n}{2}(\frac{1}{p_1}-\frac{1}{p_*})}, T_2^{1-\frac{n}{2r}}) < T_2^{-\frac{n}{2p_*}},$$

then, due to (6.2), for any $q \in [p_*, \infty]$, we get

$$\|F(u(t))\|_{L^q} \leq \tilde{C}MT_2^{-\frac{n}{2p_*}} \leq \tilde{C}KT_2^{-\frac{n}{2p_*}}, \quad t \leq 2T_2, \quad (6.9)$$

where $\tilde{C} > 0$ depends on n . Since $T_2 \geq 1$, gathering (6.8) and (6.9), for any $q \in [p_*, \infty]$, we obtain

$$\|u(t)\|_{L^q} \leq \tilde{C} \left(t^{-\frac{n}{2}(\frac{1}{p_*}-\frac{1}{q})} + T_2^{-\frac{n}{2p_*}} \right) K \leq \tilde{C} \left(t^{-\frac{n}{2}(\frac{1}{p_*}-\frac{1}{q})} + T_2^{-\frac{n}{2}(\frac{1}{p_*}-\frac{1}{q})} \right) K, \quad t \leq 2T_2.$$

Since $(2T_2)/t \geq 1$ we also get

$$\|u(t)\|_{L^q} \leq \tilde{C}t^{-\frac{n}{2}(\frac{1}{p_*}-\frac{1}{q})} K, \quad t \leq 2T_2$$

for \tilde{C} depending on n . This implies (6.7). \square

In order to prove estimate (2.6) for all times $t > 0$ we introduce once again a recursive sequence which, by uniqueness, converges to the solution u . Let

$$\begin{cases} u_0 = e^{t\Delta}\varphi \\ u_{j+1} = u_0 + \int_0^t e^{(t-s)\Delta} f(u_j(s)) ds, \quad j \geq 0. \end{cases} \quad (6.10)$$

Then, for $\varphi \geq 0$, u_j is an increasing sequence, namely,

$$u_j(t, x) \leq u_{j+1}(t, x) \leq u(t, x), \quad t > 0, \quad x \in \mathbb{R}^n,$$

for all $j \in \mathbb{N}_0$. So, if u is a solution of the Cauchy problem (1.1) and satisfies (1.8), then, for all $j \in \mathbb{N}$, u_j also satisfies

$$\sup_{t>0} \|u_j(t)\|_{\text{exp}L^2} \leq 2\|\varphi\|_{\text{exp}L^2}. \quad (6.11)$$

We end up by proving now Theorem 2.2.

Proof of Theorem 2.2. We are going to prove the first estimate (2.6) uniformly for all $u_j(t, x)$ defined as in (6.10) and so the estimate will pass to the limit $u(t, x)$. Let T_* be a sufficiently large constant to be chosen later, which satisfies

$$T_* \geq T_1 \geq 1. \quad (6.12)$$

Here T_1 is the constant given in Lemma 6.1, and depends only on K and p_* . First of all let us consider $u_0 = e^{t\Delta}\varphi$. For any $q \in [p_*, \infty]$, we have

$$\|u_0(t)\|_{L^q} \leq c_2 t^{-\frac{n}{2}\left(\frac{1}{p_*} - \frac{1}{q}\right)} \|\varphi\|_{L^{p_*}}, \quad t > 0. \quad (6.13)$$

By the monotonicity of $\{u_j\}$ and Lemma 6.2 with $T_2 = T_*$ we also have

$$\|u_j(t)\|_{L^q} \leq \tilde{C} t^{-\frac{n}{2}\left(\frac{1}{p_*} - \frac{1}{q}\right)} K, \quad 0 < t \leq 2T_*, \quad (6.14)$$

for all $j \in \mathbb{N}$. Let us assume that, for $j = m$,

$$\|u_m(t)\|_q \leq 2C_* t^{-\frac{n}{2}\left(\frac{1}{p_*} - \frac{1}{q}\right)} K, \quad t > 0, \quad (6.15)$$

with

$$C_* = \max\{c_2, \tilde{C}\},$$

and let us prove it for $j = m + 1$. Since

$$u_{m+1}(t) = u_0(t) + \int_0^t e^{(t-s)\Delta} f(u_m(s)) ds$$

we consider only the nonlinear term. For $q \in [p_*, \infty]$ and for $t > 2T_*$, we split

$$\begin{aligned} & \left\| \int_0^t e^{(t-s)\Delta} f(u_m(s)) ds \right\|_{L^q} \\ & \leq \int_0^{t/2} \left\| e^{(t-s)\Delta} f(u_m(s)) \right\|_{L^q} ds + \int_{t/2}^t \left\| e^{(t-s)\Delta} f(u_m(s)) \right\|_{L^q} ds \\ & =: J_1(t) + J_2(t). \end{aligned} \quad (6.16)$$

For $t > 2T_*$ we have

$$J_1(t) \leq c_2 \left(\int_0^{T_*} + \int_{T_*}^{t/2} \right) (t-s)^{-\frac{n}{2}\left(\frac{1}{p_1} - \frac{1}{q}\right)} \|f(u_m(s))\|_{L^{p_1}} ds := A(t) + B(t), \quad (6.17)$$

where p_1 is the constant given in (2.8). Due to estimate (6.11) we can apply Lemma 3.4 to the $A(t)$ term, and we obtain

$$\begin{aligned} A(t) &\leq Ct^{-\frac{n}{2}\left(\frac{1}{p_1}-\frac{1}{q}\right)} \int_0^{T_*} M^{1+\frac{4}{n}} ds \\ &\leq Ct^{-\frac{n}{2}\left(\frac{1}{p_1}-\frac{1}{q}\right)} T_* M^{1+\frac{4}{n}} = Ct^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)} t^{-\frac{n}{2}\left(\frac{1}{p_1}-\frac{1}{p_*}\right)} T_* M^{1+\frac{4}{n}}. \end{aligned} \quad (6.18)$$

As for the $B(t)$ term, we apply Lemma 6.1. Since it follows from (2.4), (2.8) and (6.4) with $p \in [1, 2)$ that

$$\begin{cases} p_1 = p_2 = 1 & \text{for } 1 \leq n \leq 4, \\ p_2 = \max\left\{\frac{p_* n}{n+4}, 1\right\} < \frac{2n}{n+4} = p_1 & \text{for } n \geq 5, \end{cases}$$

by (6.3) we have

$$\begin{aligned} B(t) &\leq Ct^{-\frac{n}{2}\left(\frac{1}{p_1}-\frac{1}{q}\right)} \int_{T_*}^{t/2} K^{1+\frac{4}{n}} s^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{p_1}\right)-\frac{2}{p_*}} ds \\ &\leq CK^{1+\frac{4}{n}} t^{-\frac{n}{2}\left(\frac{1}{p_1}-\frac{1}{q}\right)} \int_{T_*}^{t/2} s^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{p_1}\right)-\frac{2}{p_*}} ds. \end{aligned}$$

For $p \in (p_1, 2)$ (which implies $p_* = p$), we can choose $\sigma \in (0, 1)$ satisfying

$$0 < \sigma < \min\left\{\frac{2}{p_*} - 1, \frac{n}{2}\left(\frac{1}{p_1} - \frac{1}{p_*}\right)\right\}. \quad (6.19)$$

So, we can write

$$\begin{aligned} B(t) &\leq CK^{1+\frac{4}{n}} t^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)-\sigma} \int_{T_*}^{t/2} t^{-\frac{n}{2}\left(\frac{1}{p_1}-\frac{1}{p_*}\right)+\sigma} s^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{p_1}\right)-\frac{2}{p_*}} ds \\ &\leq CK^{1+\frac{4}{n}} t^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)-\sigma} \int_{T_*}^{t/2} s^{-\frac{n}{2}\left(\frac{1}{p_1}-\frac{1}{p_*}\right)+\sigma} s^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{p_1}\right)-\frac{2}{p_*}} ds \\ &\leq CK^{1+\frac{4}{n}} t^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)-\sigma} \int_{T_*}^{\infty} s^{-\frac{2}{p_*}+\sigma} ds \end{aligned}$$

and in the end

$$B(t) \leq CK^{1+\frac{4}{n}} t^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)-\sigma} T_*^{-\frac{2}{p_*}+\sigma+1},$$

where C is a constant, independent of M , K and T_* . For $p \leq p_1$, namely $p_* = p_1$, we cannot exploit the better decreasing in time as in (6.19). Since $p_* < 2$, for $p \leq p_1$, we get

$$B(t) \leq CK^{1+\frac{4}{n}} t^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)} T_*^{-\frac{2}{p_*}+1}. \quad (6.20)$$

In the end, for $p \in (p_1, 2)$ we have the following estimates:

$$J_1(t) \leq C \left(t^{-\frac{n}{2}\left(\frac{1}{p_1}-\frac{1}{q}\right)} T_* M^{1+\frac{4}{n}} + K^{1+\frac{4}{n}} t^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)-\sigma} T_*^{-\frac{2}{p_*}+\sigma+1} \right),$$

where C is a constant, independent of M , K and T_* . Since $2/p_* - 1 - \sigma > 0$, we can take a sufficiently large constant $T_* \geq 1$ so that

$$CK^{\frac{4}{n}}T_*^{-\frac{2}{p_*}+\sigma+1} \leq \frac{1}{4}C_*$$

which means

$$T_* \geq \left(\frac{4CK^{\frac{4}{n}}}{C_*} \right)^{\frac{1}{\frac{2}{p_*}-1-\sigma}}. \quad (6.21)$$

This together with (6.12) implies that T_* depends on K and p but not on M . Then we can also take a sufficiently small constant M so that

$$CT_*M^{\frac{4}{n}} \leq \frac{1}{4}C_*$$

and this means

$$M \leq \left(\frac{4CT_*}{C_*} \right)^{-\frac{n}{4}}. \quad (6.22)$$

In the end, since $T_* \geq 1$, for any $p \in (p_1, 2)$, by (6.1) we have

$$J_1(t) \leq C_* \left(\frac{1}{4}t^{-\frac{n}{2}\left(\frac{1}{p_1}-\frac{1}{q}\right)}M + \frac{1}{4}t^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)-\sigma}K \right) \leq \frac{C_*}{2}t^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)}K \quad (6.23)$$

for all $t > 2T_*$. In the $p \leq p_1$ case we get from (6.17), (6.18) and (6.20) similar conditions on T_* and on M in order to obtain (6.23).

Let us come back to the $J_2(t)$ term in (6.16). Since $q \geq p_* \geq p_2$, by Lemma 6.1 again, we have

$$\begin{aligned} J_2(t) &\leq \int_{t/2}^t \|f(u_j(s))\|_{L^q} ds \\ &\leq C \int_{t/2}^t K^{1+\frac{4}{n}}s^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)-\frac{2}{p_*}} ds \leq CK^{1+\frac{4}{n}}t^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)-\frac{2}{p_*}+1} \end{aligned}$$

Once again, since $p_* < 2$, we can choose $\lambda > 0$ satisfying

$$0 < \lambda < \frac{2}{p_*} - 1,$$

and we get

$$J_2(t) \leq CK^{1+\frac{4}{n}}t^{-\frac{n}{2}\left(\frac{1}{p_*}-\frac{1}{q}\right)}t^{-\frac{2}{p_*}+1+\lambda}T_*^{-\lambda}, \quad t > 2T_*,$$

where C is a constant, independent of M , K and T_* . For

$$CK^{\frac{4}{n}}T_*^{-\lambda} \leq \frac{1}{4}C_*,$$

which is implied for example by

$$T_* \geq \left(\frac{4CK^{\frac{4}{n}}}{C_*} \right)^{\frac{1}{\lambda}}, \quad (6.24)$$

we get for $t > 2T_*$

$$J_2(t) \leq \frac{C_*}{2} t^{-\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} t^{-\frac{2}{p_*} + 1 + \lambda} K. \quad (6.25)$$

This together with $T_* \geq 1$ yields

$$J_2(t) \leq \frac{C_*}{2} t^{-\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} K, \quad t \geq 2T_*. \quad (6.26)$$

Collecting (6.1), (6.13), (6.23) and (6.26) we have

$$\begin{aligned} \|u_{m+1}(t)\|_{L^q} &\leq c_2 t^{-\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} \|\varphi\|_{L^{p_*}} + C_* t^{-\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} K \\ &\leq 2C_* t^{-\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} K, \quad t > 2T_*. \end{aligned}$$

This together with (6.14) yields (6.15) with $j = m + 1$. In order to make clear the dependence of the choice we made on T_* and M , we collect below all the conditions (6.12), (6.21), (6.22), (6.24)

$$\begin{aligned} T_* &\geq \max(1, T_1), \\ T_* &\geq \left(\frac{4CK \frac{4}{n}}{C_*} \right)^{\frac{1}{\frac{2}{p_*} - 1 - \sigma}}, \\ T_* &\geq \left(\frac{4CK \frac{4}{n}}{C_*} \right)^{\frac{1}{\lambda}}, \\ M &\leq \left(\frac{4CT_*}{C_*} \right)^{-\frac{n}{4}}, \end{aligned}$$

with T_1 satisfying (6.6)

$$T_1 \geq \left(\frac{(CK)^2}{\log 2} \right)^{\frac{p_*}{n}},$$

and C_* , C , η , σ and λ constants depending at most on n and p_* . In the end, we find a function F depending on n , p_* , K such that the condition on M can be written as

$$M \leq F(n, p_*, K)$$

as it was announced in the statement of Theorem 2.2. Finally we prove (2.7). By (6.23) and (6.25), for $p \in (p_1, 2)$, we have

$$t^{\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} \|u(t) - e^{t\Delta} \varphi\|_{L^q} = o(1), \quad t \rightarrow \infty.$$

Now, by density, let $\{\varphi_n\} \subset C_0^\infty$ such that $\varphi_n \rightarrow \varphi$ in L^{p_*} . Then

$$\begin{aligned} t^{\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} \|e^{t\Delta} \varphi\|_{L^q} &\leq t^{\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} \|e^{t\Delta} (\varphi - \varphi_n)\|_{L^q} + t^{\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} \|e^{t\Delta} \varphi_n\|_{L^q} \\ &\leq c_2 \left(\|\varphi - \varphi_n\|_{L^{p_*}} + t^{\frac{n}{2} \left(\frac{1}{p_*} - \frac{1}{q} \right)} t^{-\frac{n}{2} \left(1 - \frac{1}{q} \right)} \|\varphi_n\|_{L^1} \right) \\ &\leq c_2 \left(\|\varphi - \varphi_n\|_{L^{p_*}} + t^{-\frac{n}{2} \left(1 - \frac{1}{p_*} \right)} \|\varphi_n\|_{L^1} \right) \end{aligned}$$

for all $t > 0$. This proves that

$$t^{\frac{n}{2}\left(\frac{1}{p^*}-\frac{1}{q}\right)}\|e^{t\Delta}\varphi\|_{L^q} = o(1), \quad t \rightarrow \infty$$

and so

$$t^{\frac{n}{2}\left(\frac{1}{p^*}-\frac{1}{q}\right)}\|u(t)\|_{L^q} = o(1), \quad t \rightarrow \infty.$$

Thus the proof of Theorem 2.2 is complete. \square

Remark 6.3 *It is worth commenting on the meaning of condition (2.5) appearing in Theorem 2.2 about the "smallness" of the $\exp L^2$ norm of the initial data φ . In fact, the evolution equation governing the Cauchy problem (1.1) has no scaling invariance and the L^p and $\exp L^2$ norms have no relationship between each other. In order to have initial data which fulfill condition (2.5), let us choose a function $\varphi \in L^p \cap L^\infty$ with $p \in [1, 2)$. Since $\varphi \in L^2 \cap L^\infty$, then $\varphi \in \exp L^2$ (see [1]). Then, let us consider a dilation $\varphi_\lambda(x) = \lambda^{\frac{n}{p}}\varphi(\lambda x)$ so that $\|\varphi_\lambda\|_{L^p} = \|\varphi\|_{L^p}$. Since $\|\varphi_\lambda\|_{L^2} = \lambda^{n\left(\frac{1}{p}-\frac{1}{2}\right)}\|\varphi\|_{L^2}$ and $\|\varphi_\lambda\|_{L^\infty} = \lambda^{\frac{n}{p}}\|\varphi\|_{L^\infty}$, it follows*

$$\limsup_{\lambda \rightarrow 0} \|\varphi_\lambda\|_{\exp L^2} \leq \lim_{\lambda \rightarrow 0} (\|\varphi_\lambda\|_{L^2} + \|\varphi_\lambda\|_{L^\infty}) = 0.$$

This implies that there is $\lambda > 0$ so that φ_λ fulfills condition (2.5), even though its L^p norm might be large.

We end this section by proving Theorem 2.3. In the following Lemma we assume $\|v(s)\|_{L^q}$ bounded at the origin and decaying at infinity, and we can deduce that also $\|f(v(s))\|_{L^r}$ is bounded and decays at infinity for $r \geq p_2$ where p_2 is defined in (6.4).

Lemma 6.3 *Let $n \geq 1$, $p \in [1, 2)$ and $L > 0$. Let $v(t, x)$ a function satisfying*

$$\sup_{s>0} (1+s)^{\frac{n}{2}\left(\frac{1}{p}-\frac{1}{q}\right)}\|v(s)\|_{L^q} \leq CL \tag{6.27}$$

for all $q \in [p, \infty]$ and C independent of q . Then, there is $\delta > 0$ such that if $\tilde{K} < \delta$, then for all $r \in [p_2, \infty]$

$$\sup_{s>0} (1+s)^{\frac{n}{2}\left(\frac{1}{p}-\frac{1}{r}\right)+\frac{2}{p}}\|f(v(s))\|_{L^r} \leq 2(CL)^{1+\frac{4}{n}} \tag{6.28}$$

where f and p_2 are defined as in (1.6) and (6.4), respectively.

Proof. Let $r_1 = 4/n + 1$. Then, since $r_1 p_2 \geq p$, for any $r \in [p_2, \infty]$, it follows from (1.6),

(3.4) and (6.27) that

$$\begin{aligned}
\|f(v(s))\|_{L^r} &\leq \sum_{k=0}^{\infty} \frac{\|v^{2k+r_1}(s)\|_{L^r}}{k!} \\
&\leq \sum_{k=0}^{\infty} \frac{\|v(s)\|_{L^{(2k+r_1)r}}^{2k+r_1}}{k!} \\
&\leq \sum_{k=0}^{\infty} \frac{1}{k!} \left(C(1+s)^{-\frac{n}{2} \left(\frac{1}{p} - \frac{1}{(2k+r_1)r} \right)} L \right)^{2k+r_1} \\
&\leq (C\tilde{K})^{r_1} (1+s)^{\frac{n}{2r} - \frac{nr_1}{2p}} \sum_{k=0}^{\infty} \frac{1}{k!} \left(C(1+s)^{-\frac{n}{2p}} L \right)^{2k} \\
&\leq (C\tilde{K})^{r_1} (1+s)^{\frac{n}{2r} - \frac{nr_1}{2p}} \sum_{k=0}^{\infty} \frac{1}{k!} (CL)^{2k}
\end{aligned} \tag{6.29}$$

for all $s > 0$. We can take a sufficiently small δ , which independent of r , so that, for $L \leq \delta$, it holds that

$$\sum_{k=0}^{\infty} \frac{1}{k!} (CL)^{2k} = e^{(C\tilde{K})^2} \leq 2.$$

This together with (6.29) implies (6.28) Thus Lemma 6.3 follows. \square

Proof of Theorem 2.3. Put $L = \max\{\|\varphi\|_{\exp L^2}, \|\varphi\|_{L^p}\}$. Applying the same argument as in the proof of Theorem 2.1 with Lemma 6.3, we can prove Theorem 2.3. So we omit the proof. \square

7 Asymptotic behavior

Let us come to the asymptotic behavior of the solution u as stated in Theorem 2.4.

Proof of Theorem 2.4. By the assumption for the initial data φ with (1.8) we can apply Lemma 3.4 and obtain

$$\sup_{t>0} \|f(u(t))\|_{L^1} < \infty, \quad 1 \leq n \leq 4.$$

For the case $n \geq 5$, since

$$f(u(t)) = u^{1+\frac{4}{n}}(t) e^{u^2(t)} = u^{1+\frac{4}{n}}(t) + u^{1+\frac{4}{n}}(t) \left(e^{u^2(t)} - 1 \right),$$

by a proof analogous to Lemma 3.4 and (2.10) we get

$$\sup_{t>0} \|f(u(t))\|_{L^1} \leq \sup_{t>0} \left(\|u(t)\|_{L^{1+\frac{4}{n}}}^{1+\frac{4}{n}} + \left\| u^{1+\frac{4}{n}}(t) \left(e^{u^2(t)} - 1 \right) \right\|_{L^1} \right) < \infty.$$

Therefore we can define a mass of $u(t)$ denote by $m(t)$, that is,

$$m(t) := \int_{\mathbb{R}^n} \varphi(x) dx + \int_0^t \int_{\mathbb{R}^n} f(u(s, x)) dx ds, \quad t \geq 0.$$

In fact, if $\varphi \in L^1$, then we can define the integral equation (1.7) in L^1 for almost all $t > 0$, and by (G1) we have

$$\begin{aligned} \int_{\mathbb{R}^n} e^{t\Delta} \varphi(x) dx &= \int_{\mathbb{R}^n} \varphi(x) dx, \\ \int_{\mathbb{R}^n} \int_0^t e^{(t-s)\Delta} f(u(s, x)) ds dx &= \int_0^t \int_{\mathbb{R}^n} f(u(s, x)) dx ds. \end{aligned}$$

Let T_1 be the constant given in Lemma 6.1. Then, by Lemma 6.1 we have

$$\begin{aligned} \int_0^t \int_{\mathbb{R}^n} f(u(s, x)) dx ds &= \int_0^{T_1} \int_{\mathbb{R}^n} f(u(s, x)) dx ds + \int_{T_1}^t \int_{\mathbb{R}^n} f(u(s, x)) dx ds \\ &\leq CT_1 + C \int_{T_1}^t s^{-2} ds \leq C(T_1 + T_1^{-1}) \end{aligned} \quad (7.1)$$

for all $t \geq T_1$. This implies that there exists the limit of $m(t)$, which we denote by m_* , such that

$$m_* := \lim_{t \rightarrow \infty} m(t) = \int_{\mathbb{R}^n} \varphi(x) dx + \int_0^\infty \int_{\mathbb{R}^n} f(u(s, x)) dx ds.$$

Furthermore, similarly to (7.1), we obtain

$$m_* - m(t) \leq C \int_t^\infty \int_{\mathbb{R}^n} f(u(s, x)) dx ds \leq Ct^{-1}$$

for all $t \geq T_1$. Therefore, applying an argument similar to the proof of Theorem 1.1 in [5] with (2.9) and (2.10), we have (2.11), and the proof of Theorem 2.4 is complete. \square

8 Generalization

In this last section, we are going to get similar results as in the Section 2 to the general problem. Let $\theta \in (0, 2]$ and $r > 1$. Put

$$f(u) = |u|^{\frac{r\theta}{n}} u e^{u^r}. \quad (8.1)$$

Then, similarly to Lemma 3.4, the following hold.

Lemma 8.1 *Let $n \geq 1$ and $M > 0$. Assume that a function $u \in L^\infty(0, \infty; \exp L^r)$ satisfies the condition*

$$\sup_{t>0} \|u(t)\|_{\exp L^r} \leq M.$$

Let f be the function defined as in (8.1). Put

$$p_1 = \max\{rn/(n + r\theta), 1\}.$$

Then, for all $p \in [p_1, \infty)$ there is $\varepsilon = \varepsilon(p) > 0$ such that if $M < \varepsilon$, then

$$\sup_{t>0} \|f(u(t))\|_{L^p} \leq 2Cp^3 M^{1+\frac{r\theta}{n}},$$

where C is independent of p , n and M .

Let us consider now for simplicity only the integral equation

$$u(t) = e^{-t\mathcal{L}_\theta} \varphi + \int_0^t e^{-(t-s)\mathcal{L}_\theta} f(u(s)) ds. \quad (8.2)$$

This is the integral formulation of (2.12). In a similar way as in Section 4 one can prove that this integral equation is equivalent to the differential equation if we consider small solutions.

Applying the arguments as in the previous sections with Lemma 8.1, we have the following.

Theorem 8.1 *Let $n \geq 1$ and $r > 1$. Assume $\varphi \in \exp L^r$. Then there exists $\varepsilon = \varepsilon(n) > 0$ such that, if $\|\varphi\|_{\exp L^r} < \varepsilon$, then there exists a unique solution u of the integral equation (8.2) satisfying*

$$u \in L^\infty(0, \infty; \exp L^r)$$

and

$$\sup_{t>0} \|u(t)\|_{\exp L^r} \leq 2\|\varphi\|_{\exp L^r}. \quad (8.3)$$

Theorem 8.2 *Let $n \geq 1$, $r > 1$ and $\varphi \in \exp L^r$ with $\varphi \geq 0$. Assume that there exists a unique positive solution u of (8.2) satisfying (8.3). Then there exist $\varepsilon = \varepsilon(n) > 0$ and $C = C(n) > 0$ such that, if $\|\varphi\|_{\exp L^r} < \varepsilon$, then the solution u satisfies*

$$\|u(t)\|_{L^q} \leq Ct^{-\frac{n}{\theta} \left(\frac{1}{r} - \frac{1}{q} \right)} \|\varphi\|_{\exp L^r}, \quad t > 0,$$

for any $q \in [r, \infty]$.

Theorem 8.3 *Assume the same conditions as in Theorem 8.2. Furthermore, suppose that $\varphi \in L^p$ for some $p \in [1, r)$. Put*

$$p^* = \max \left\{ p, \frac{rn}{n + r\theta} \right\}.$$

Then there exist positive constants $\varepsilon = \varepsilon(n)$, $C = C(n)$ and a positive function $F = F(n, p^, \|\varphi\|_{L^{p^*}})$ such that, if*

$$\|\varphi\|_{\exp L^r} < \min(\varepsilon, F(n, p^*, \|\varphi\|_{L^{p^*}})),$$

then the solution u satisfies

$$\|u(t)\|_{L^q} \leq Ct^{-\frac{n}{\theta} \left(\frac{1}{p^*} - \frac{1}{q} \right)} \|\varphi\|_{\exp L^r \cap L^{p^*}}, \quad t > 0, \quad (8.4)$$

for any $q \in [p^, \infty]$. In particular, if $p \in (p_3, r)$, then*

$$\|u(t)\|_{L^q} = o \left(t^{-\frac{n}{\theta} \left(\frac{1}{p^*} - \frac{1}{q} \right)} \right), \quad t \rightarrow \infty. \quad (8.5)$$

Here $[K]$ is the integer satisfying $K - 1 \leq [K] < K$ and

$$p_3 := \max \left\{ 1, \frac{rn}{n + r\theta} \right\}.$$

Theorem 8.4 *Assume the same conditions as in Theorem 8.3. Then there exists a positive constant $\delta = \delta(n)$ such that, if*

$$\max\{\|\varphi\|_{\exp L^r}, \|\varphi\|_{L^p}\} < \delta,$$

then (8.4) with $p^ = p$ holds for all $q \in [p, \infty]$. In particular, for all $q \in [p, \infty)$,*

$$\|u(t)\|_{L^q} \leq C(1+t)^{-\frac{n}{\theta}\left(\frac{1}{p}-\frac{1}{q}\right)} \|\varphi\|_{\exp L^r \cap L^p}, \quad t > 0.$$

Furthermore, if $p \in (1, r)$, then (8.5) with $p^ = p$ holds.*

Theorem 8.5 *Let $n \geq 1$, $\varphi \geq 0$ and $\varphi \in \exp L^r \cap L^1$. Assume $\|\varphi\|_{\exp L^r}$ is small enough. Furthermore, suppose that*

a) for $n \geq 1$,

$$\sup_{t>0} t^{\frac{n}{\theta}\left(1-\frac{1}{q}\right)} \|u(t)\|_{L^q} < \infty, \quad q \in [1, \infty];$$

b) for $n > \max\{1, [r\theta/(r-1)]\}$, assume moreover that there is $T^ > 0$ such that*

$$\sup_{0<t<T^*} \|u(t)\|_{L^{\frac{r\theta}{n}+1}} < \infty.$$

Then there exists the limit

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}^n} u(x, t) dx = \int_{\mathbb{R}^n} \varphi(x) dx + \int_0^\infty \int_{\mathbb{R}^n} f(u(t, x)) dx dt := m_*$$

such that

$$\lim_{t \rightarrow \infty} t^{\frac{n}{\theta}\left(1-\frac{1}{q}\right)} \|u(t) - m_* G_\theta(t+1)\|_{L^q} = 0,$$

for any $q \in [1, \infty]$.

Acknowledgements. The work of the second author (T. Kawakami) was supported in part by Grant-in-Aid for Young Scientists (B) (No. 24740107) and (No. 16K17629) of JSPS (Japan Society for the Promotion of Science) and by the JSPS Program for Advancing Strategic International Networks to Accelerate the Circulation of Talented Researchers “Mathematical Science of Symmetry, Topology and Moduli, Evolution of International Research Network based on OCAMI”.

References

- [1] R. A. Adams and J. J. F. Fournier, *Sobolev Spaces, 2nd. ed.*, Academic Press, New York, 2003.
- [2] H. Brezis and T. Cazenave, A nonlinear heat equation with singular initial data, *J. Anal. Math.* **68** (1996) 277–304.

- [3] T. Cazenave and F. B. Weissler, The Cauchy problem for the critical nonlinear Schrödinger equation in H^s , *Nonlinear Anal.* **14** (10) (1990) 807–836.
- [4] M. Escobedo and O. Kavian, Variational problems related to self-similar solutions of the heat equation, *Nonlinear Anal.* **11** (1987), 1103–1133.
- [5] M. Fila, K. Ishige and T. Kawakami, Convergence to the Poisson kernel for the Laplace equation with a nonlinear dynamical boundary condition, *Commun. Pure Appl. Anal.* **11** (3) (2012) 1285–1301.
- [6] H. Fujita, On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$, *J. Fac. Sci. Univ. Tokyo Sect. I* **13** (1966) 109–124.
- [7] A. Haraux and F. B. Weissler, Non-uniqueness for a semilinear initial value problem, *Indiana Univ. Math. J.* **31** (1982) 167–189.
- [8] N. Ioku, The Cauchy problem for heat equations with exponential nonlinearity, *J. Differential Equations* **251** (4-5) (2011) 1172–1194.
- [9] N. Ioku, Regularity theory of exponential type for elliptic and parabolic partial differential equations in Orlicz spaces, PhD thesis, Tohoku University, Sendai, Japan, 2011.
- [10] N. Ioku, B. Ruf and E. Terraneo, Existence, non-existence, and uniqueness for a heat equation with exponential nonlinearity in \mathbb{R}^2 , *Math. Phys. Anal. Geom.* **18** (1) (2015) Art. 29, 19 pp.
- [11] K. Ishige, T. Kawakami and K. Kobayashi, Asymptotics for a nonlinear integral equation with a generalized heat kernel, *J. Evol. Equ.* **14** (2014) 749–777.
- [12] K. Ishige, T. Kawakami and K. Kobayashi, Global solutions for a nonlinear integral equation with a generalized heat kernel, *Discrete Contin. Dyn. Syst. Ser. S* **7** (2014) 767–783.
- [13] T. Kawanago, Existence and behaviour of solutions for $u_t = \Delta(u^m) + u^l$, *Adv. Math. Sci. Appl.* **7** (1997) 367–400.
- [14] T. Y. Lee and W. M. Ni, Global existence, large time behavior and life span of solutions of a semilinear parabolic Cauchy problem, *Trans. Amer. Math. Soc.* **333** (1992) 365–378.
- [15] M. Nakamura and T. Ozawa, Nonlinear Schrödinger equations in the Sobolev space of critical order, *J. Funct. Anal.* **155** (2) (1998) 364–380.
- [16] P. Quittner and P. Souplet, *Superlinear parabolic problems: Blow-up, global existence and steady states*, Birkhäuser Advanced Texts, Basel, 2007.
- [17] M. M. Rao and Z. D. Ren, *Theory of Orlicz Spaces*, Marcel Dekker, Inc., New York, 1991.
- [18] F. Ribaud, Cauchy problem for semilinear parabolic equations with initial data in $H_p^s(\mathbb{R}^n)$ spaces, *Rev. Mat. Iberoamericana* **14** (1998) 1–46.

- [19] B. Ruf and E. Terraneo, The Cauchy problem for a semilinear heat equation with singular initial data, *Progr. Nonlinear Differential Equations Appl.* **50** (2002) 295–309.
- [20] N. S. Trudinger, On imbeddings into Orlicz spaces and some applications, *J. Math. Mech.* **17** (1967) 473–483.
- [21] X. Wang, On the Cauchy problem for reaction-diffusion equations, *Trans. Amer. Math. Soc.* **337** (2) (1993) 549–590.
- [22] F. B. Weissler, Local existence and nonexistence for semilinear parabolic equations in L^p . *Indiana Univ. Math. J.* **29** (1980) 79–102.
- [23] F. B. Weissler, Existence and nonexistence of global solutions for a semilinear heat equation, *Israel J. Math.* **38** (1981) 29–40.

Addresses:

Giulia Furioli
 DIGIP, Università di Bergamo,
 Viale Marconi 5, I-24044 Dalmine (BG), Italy
 E-mail: gfurtoli@unibg.it

Tatsuki Kawakami
 Department of Mathematical Sciences,
 Osaka Prefecture University,
 Gakuencho 1-1, Sakai 599-8531, Japan
 E-mail: kawakami@ms.osakafu-u.ac.jp

Bernhard Ruf
 Dipartimento di Matematica F. Enriques,
 Università degli studi di Milano,
 Via Saldini 50 , I-20133 Milano, Italy
 E-mail: bernhard.ruf@unimi.it

Elide Terraneo
 Dipartimento di Matematica F. Enriques,
 Università degli studi di Milano,
 Via Saldini 50 , I-20133 Milano, Italy
 E-mail: elide.terraneo@unimi.it